

# Pharmacy Management Drug Policy

**SUBJECT:** Growth Hormone Policy  
**POLICY NUMBER:** Pharmacy-18  
**EFFECTIVE DATE:** 08/03  
**LAST REVIEW DATE:** 3/6/2019

*If the member's subscriber contract excludes coverage for a specific service or prescription drug, it is not covered under that contract. In such cases, medical or drug policy criteria are not applied. Medical or drug policies apply to commercial and Health Care Reform products only when a contract benefit for the specific service exists.*

## **DESCRIPTION:**

Human growth hormone (GH), also known as somatotropin, is synthesized in the somatotrophs of the anterior pituitary gland. Synthetic growth hormone (trade names: Genotropin, Humatrope, Norditropin, Nutropin AQ, Omnitrope, Saizen, Serostim, Zomacton, Zorbtive) is primarily used as replacement therapy in patients with growth hormone deficiency (GHD). It is also used as anabolic therapy in patients with the wasting syndrome of AIDS, patients suffering malnourishment due to short bowel syndrome and patients suffering from third degree burns. Zorbtive is approved for treatment of short bowel syndrome in patients experiencing malabsorption, malnutrition, weight loss, or dehydration.

## **POLICY:**

Based on our assessment and review of peer-reviewed literature, the administration of human growth hormone (GH) has been proven to be effective and therefore, **medically necessary** for the following conditions:

### **I. CHILDREN - INCLUSION CRITERIA**

Only a pediatric endocrinologist should prescribe GH for children. In children with renal insufficiency, GH therapy can be managed by a pediatric nephrologist with expertise in growth hormone therapy.

<b><u>If the FDA approved indication is:</u></b>	<b><u>Then the criteria for review includes the following for medical necessary consideration:</u></b>
Growth Hormone Deficiency (GHD) diagnosis in children	<ul style="list-style-type: none"> <li>• Requires the following criteria (<b>#1 OR #2</b>):</li> <li>1. Diminished Growth Hormone Response (level less than 10 ng/ml) to <b>2 or more</b> of the following provocative tests:            *Note – must be 2 <i>different</i> agents           <ul style="list-style-type: none"> <li>• Levodopa,</li> <li>• Insulin-induced hypoglycemia,</li> <li>• Arginine,</li> <li>• Clonidine,</li> <li>• Glucagon</li> </ul> </li> <li><b>OR</b></li> <li>2. Low IGF-1 (insulin-like growth factor) for age, sex, and pubertal status in children age 6 or greater in the absence of chronic disease (such as, malnutrition, hepatic disease, renal insufficiency, diabetes, and hypothyroidism) in combination with height velocity (HV) less than 25th percentile in 6-12 months prior to GH therapy</li> </ul>

## Pharmacy Management Drug Policy

### Growth Hormone

	<p>In the instance of discrepancy between IGF-1 &amp; HV (i.e., IGF-1 is normal &amp; HV &lt;25th percentile) growth hormone stimulation testing (#1) will be required.</p> <p><b>AND at least 2 of the following:</b></p> <ol style="list-style-type: none"> <li>Growth velocity less than 7 cm/year before age 3 years, <b>or</b> less than 4-5 cm/year from age 3 years to onset of puberty. Severe short stature is defined as a height more than 2 standard deviations (SD) below the population mean.</li> <li>Delayed bone age - greater than 2 S.D. below mean for chronological age, generally greater than 2 years delayed in patients with radiographic evidence of <i>epiphyses not closed</i>.</li> <li>A known risk factor for Growth Hormone Deficiency such as craniofacial anomalies, central nervous system structural abnormalities, congenital hypopituitarism, panhypopituitarism, or syndromes associated with hypopituitarism, or children who have had hypophysectomy (surgical removal of pituitary gland) or history of central nervous system irradiation, including children who undergone brain radiation.</li> </ol>
Turner Syndrome	<ul style="list-style-type: none"> <li>Treatment of short stature in Turner Syndrome defined as a 45, XO genotype or mosaic 45XO 46 XY. Treatment should be initiated, as soon as patients fall below the 5<sup>th</sup> percentile of the normal growth curve for girls but not younger than age 2 years old.</li> </ul>
Short Stature <b>WITH</b> renal insufficiency (CKD)	<ul style="list-style-type: none"> <li>Children with height less than 3<sup>rd</sup> percentile for chronological age with renal insufficiency defined as serum creatinine of greater than 3.0 mg/dl or a creatinine clearance between 5 and 75/ml/min per 1.73m<sup>2</sup> before renal transplant. <i>Not recommended for post-transplant patients.</i></li> </ul>
Prader-Willi Syndrome with short stature or growth failure	<ul style="list-style-type: none"> <li>Children with Prader-Willi Syndrome with short stature or growth failure. Growth hormone is contraindicated in patients with Prader-Willi syndrome who are severely obese or have severe respiratory impairment.</li> </ul>
Noonan Syndrome with short stature	<ul style="list-style-type: none"> <li>Patient's height must be greater than 2 SD below the mean for gender &amp; age.</li> </ul>
Short Stature homeobox-containing gene (SHOX)	<ul style="list-style-type: none"> <li>Children with SHOX (short stature homeobox-containing gene) deficiency demonstrated by chromosome analysis and whose epiphyses are not closed.</li> </ul>

## Pharmacy Management Drug Policy

### Growth Hormone

<p>Intrauterine growth retardation (including those with Russell – Silver syndrome) or small for gestational age (SGA)</p>	<ul style="list-style-type: none"> <li>• <b>Growth hormone is indicated</b> for short stature associated with SGA in children who did not catch up by 2 years of age. These children do not exhibit GH deficiency. All of the following criteria must be met:             <ol style="list-style-type: none"> <li>1. Patient must be evaluated by a pediatric endocrinologist <b>and</b></li> <li>2. Patient must have been born SGA. SGA is defined as birth weight of less than 2500 grams at a gestational age of more than 37 weeks or length below the 3<sup>rd</sup> percentile for gestational age or birth weight and/or length at least 2 SDs below the mean for gestational age and gender. Most children born SGA will show catch up growth by age 2. <b>and</b></li> <li>3. Age - It is recommended that therapy be initiated between the ages of 2 and 8 years. The effect of GH on SGA children is greater when GH is given to those younger than 4 years of age.                 <ul style="list-style-type: none"> <li>• Consideration for patients greater than 8 years of age will only be given if the child is prepubertal. Efficacy has not been established in pubertal adolescents born SGA.</li> </ul> <b>AND</b> <ul style="list-style-type: none"> <li>• Therapy should be discontinued when growth velocity is less than 5cm/year or evidence of epiphyseal fusion is present.</li> </ul> </li> </ol> </li> </ul>
--	---

## II. ADULTS - INCLUSION CRITERIA

GH treatment for adults must be requested and coordinated by an endocrinologist. Adults with somatotropin deficiency syndrome require **at least one** of the following criteria (**A or B**):

### A. Diagnosis confirmed by chemical documentation:

1. Insulin Tolerance Test (ITT) less than 5ng/mL. ITT is the test of choice. This test is contraindicated for patients with the following:
  - Age greater than 65 years **OR**
  - History of ischemic heart disease or cerebrovascular disease **OR**
  - Abnormal EKG **OR**
  - Seizure disorders
2. If ITT is contraindicated then the following tests may be considered - IV arginine in combination with GH-releasing hormone (GHRH) less than 9-10ng/mL
3. If there is laboratory documented deficiencies of 3 or more pituitary hormones, insulin tolerance and arginine tests are not required.
4. Serum IGF-I below normal. However, a normal IGF-I does not exclude diagnosis of GHD. This test should be used in conjunction with other diagnostic tests to determine presence of GHD. Levels of IGF-I may be reduced by poor nutrition, severe hepatic disease, poorly controlled diabetes mellitus, and inadequately treated hypothyroidism.

**OR**

## Pharmacy Management Drug Policy

### Growth Hormone

**B. GH is considered medically necessary for adult patients who meet both #1 and #2 along with one criteria from #3:**

1. Biochemical diagnosis of somatotropin deficiency syndrome, by means of a negative response to a standard growth hormone stimulation test as noted above maximum peak less than 5ng/mL (regardless of stimulation test or GH assay used, the cutoff point of 5mcg/mL is used for all provocative tests.) **AND**
2. Exhibit clinical symptoms of somatotropin deficiency syndrome such as
  - Increased weight and body fat mass, decreased lean body mass,
  - Decreased exercise capacity,
  - Decreased muscle mass and strength,
  - Reduced cardiac performance,
  - Reduced bone density and increased fracture rate, and
  - Poor sleep, impaired sense of well-being, lack of motivation.
3. **AND ONE** of the following:
  - a. Adult onset: patients with somatotropin deficiency syndrome and multiple hormone deficiencies (hypopituitarism or panhypopituitarism) as a result of:
    - Pituitary disease or
    - Hypothalamic disease or
    - Pituitary surgery or
    - Radiation therapy directly to or involving the pituitary gland.
  - b. Child Onset: patients who were growth hormone deficient during childhood and who have somatotropin deficiency confirmed as an adult
  - c. Sheehan's syndrome (pituitary infarction)
  - d. Autoimmune hypophysitis
  - e. Hypophysitis associated with other inflammatory conditions (e.g. Sarcoidosis, etc.).

### **III. CHILDREN AND ADULT INCLUSION CRITERIA**

GH therapy is medically necessary if the patient meets **one** of the following:

- A.** Patients with AIDS wasting or cachexia or children with HIV associated failure to thrive defined as a greater than 10% of baseline weight loss or weight of <90% of ideal body weight **AND**
  - Chronic diarrhea (at least 2 loose stools per day for at least 30 days) **or**
  - Chronic weakness that cannot be explained by a concurrent illness other than HIV infection. Patients must be simultaneously treated with antiviral agents. Diet must provide at least 100% of estimated caloric requirement. Evaluate weight on a quarterly basis for patients being treated for HIV wasting
  - **For recertification**, the patient must have gained at least 2 kg of body weight after 12 weeks of therapy (clinical trials averaged 3.2kg)
- B.** Patients suffering from third degree burns.
- C.** Patients with short bowel syndrome who are receiving specialized nutritional support, defined as, high carbohydrate low-fat diet that is adjusted for individual patient requirements and preferences. Patients with short bowel syndrome who are experiencing malabsorption, malnutrition, weight loss, or dehydration. Approval will be limited to a 4 week course per year. Special consideration will be given to any request for therapy longer in duration than 4 weeks.

# Pharmacy Management Drug Policy

## Growth Hormone

### **RECERTIFICATIONS**

For children with Growth Hormone Deficiency, approval will be for 12 months at a time. There must be documented improvement for patients to continue receiving growth hormone replacement. The following documentation must be submitted for review every 12 months:

- The difference between the patient's current height and their predicted adult height
  - The patient's current bone age and chronological age
  - The patient's current height percentile on height/weight chart
  - Puberty status
  - Treatment Plans/Goals
1. In children, GH therapy is typically discontinued when the growth velocity is less than 5 cm per year, **OR** when epiphyseal fusion has occurred, **OR** when the predicted mid-parental height is reached.
  2. Radiographic testing to determine if epiphyses are closed at age 14 in girls and at age 16 in boys is required.

### **POLICY GUIDELINES:**

1. Prior authorization is contract dependent.
2. **Starting June 15, 2010 all requests for growth hormone therapy will be required to use Omnitrope except in the following instances**
  - a. Serostim will be approved for wasting or cachexia associated with HIV
  - b. Zorbtive will be approved for the treatment of short bowel syndrome
  - c. Individuals who have a documented sensitivity to benzyl alcohol (a preservative in Omnitrope 5 Pen and Omnitrope 5.8mg/vial) and to phenol (a preservative in Omnitrope 10 Pen) will be authorized to use Genotropin or Humatrope (which contain a different preservative)
3. Benzyl alcohol should not be used in children under the age of 3. Omnitrope 10 should be used in children under the age of 3 as it does not contain benzyl alcohol.
4. In children with chronic renal insufficiency, GH therapy is discontinued at the time of the renal transplant. Continued growth failure after a successful renal transplant may indicate the need for re-initiation of growth hormone therapy.
5. Discontinue if growth rate is <5cm/yr.
6. Discontinue if body mass stores normalized in HIV patients.
7. This policy is applicable to drugs that are included on a specific drug formulary. If a drug referenced in this policy is non-formulary, please reference the Coverage Exception Evaluation Policy for All Lines of Business Formularies policy for review guidelines.
8. For contracts where Insurance Law § 4903(c-1), and Public Health Law § 4903(3-a) are applicable, if trial of preferred drug(s) is the only criterion that is not met for a given condition, and one of the following circumstances can be substantiated by the requesting provider, then trial of the preferred drug(s) will not be required.
  - The required prescription drug(s) is (are) contraindicated or will likely cause an adverse reaction or physical or mental harm to the member;
  - The required prescription drug is expected to be ineffective based on the known clinical history and conditions and concurrent drug regimen;

## Pharmacy Management Drug Policy

### Growth Hormone

- The required prescription drug(s) was (were) previously tried while under the current or a previous health plan, or another prescription drug or drugs in the same pharmacologic class or with the same mechanism of action was (were) previously tried and such prescription drug(s) was (were) discontinued due to lack of efficacy or effectiveness, diminished effect, or an adverse event;
  - The required prescription drug(s) is (are) not in the patient's best interest because it will likely cause a significant barrier to adherence to or compliance with the plan of care, will likely worsen a comorbid condition, or will likely decrease the ability to achieve or maintain reasonable functional ability in performing daily activities;
  - The individual is stable on the requested prescription drug. The medical profile of the individual (age, disease state, comorbidities), along with the rationale for deeming stability as it relates to standard medical practice and evidence-based practice protocols for the disease state will be taken into consideration.
  - The above criteria are not applicable to requests for brand name medications that have an AB rated generic. We can require a trial of an AB-rated generic equivalent prior to providing coverage for the equivalent brand name prescription drug.
9. Unless otherwise stated above within the individual drug criteria, approval time period will be for 2 years.
- Continued approval at time of recertification will require documentation that the drug is providing ongoing benefit to the patient in terms of improvement or stability in disease state or condition. Such documentation may include progress notes, imaging or laboratory findings, and other objective or subjective measures of benefit which support that continued use of the requested product is medically necessary. Also, ongoing use of the requested product must continue to reflect the current policy's preferred formulary. Recertification reviews may result in the requirement to try more cost-effective treatment alternatives as they become available (i.e.; generics, biosimilars, or other guideline-supported treatment options). Requested dosing must continue to be consistent with FDA-approved or off-label/guideline-supported dosing recommendations.
10. This policy is applicable to drugs that are included on a specific drug formulary. If a drug referenced in this policy is non-formulary, please reference the Coverage Exception Evaluation Policy for All Lines of Business Formularies policy for review guidelines.

#### **INVESTIGATIONAL EXCLUSIONS:**

Conditions considered **investigational** due to lack of peer-reviewed literature for which efficacy or safety data is not yet available include, but are not limited to:

- Constitutional delay of growth and development,
- Skeletal dysplasias,
- Osteogenesis Imperfecta,
- Anabolic therapy provided to counteract an acute or chronic catabolic illness (i.e., surgery outcomes, trauma, critical illnesses); except for AIDS
- Chronic Fatigue Syndrome
- Fibromyalgia
- Battered Wife Syndrome
- Obesity
- Cystic Fibrosis
- Crohn's disease
- Transplantation



## Pharmacy Management Drug Policy

### Growth Hormone

- MSA, Multiple System Atrophy
- Delay of puberty in combination with GnRH to enhance linear growth with GH

#### **EXCLUSIONS**

1. Acute Critical Illness
2. Anti-aging therapy
3. Pregnancy (not an absolute contraindication, not approved by the FDA/Infertility)
4. Hypersensitivity
5. Cardiomyopathy and heart failure
6. Idiopathic short stature and familial short stature - Pediatric patients who are non-GHD with short- stature (also known as ISS), as studies have failed to demonstrate a significant impact of height on psychosocial morbidity. The American Academy of Pediatrics (AAP) has pointed out that there will always be a population of individuals considered short based on the normal distribution of height, regardless of how the bell-shaped curve may be altered by GH therapy.
7. Down Syndrome and other syndromes associated with short stature and malignant diathesis.
8. Anabolic therapy to enhance body mass or strength for professional, recreational or social reasons.
9. GH is contraindicated in patients with an active malignant condition. If GHD results from an intracranial tumor, absence of tumor growth or tumor recurrence should be documented for 6 to 12 months before initiation of GH treatment.
10. Proliferative Diabetic Retinopathy
11. Pseudotumor Cerebri
12. Weight loss or medical weight loss programs (taken orally or injected)
13. Documented closure of epiphysis
14. GH is contraindicated for individuals with Prader-Willi syndrome who are severely obese or have severe respiratory impairment.
15. Omnitrope 5 & 5.8mg which contains benzyl alcohol as a preservative is contraindicated in children under the age of 3. Omnitrope 10 contains phenol as a preservative and is safe to use in children of all ages.

#### **For all patients covered under all Managed Medicaid Plans**

Effective 6/1/2012, NYS Medicaid will no longer provide coverage for Growth Hormone therapy for diagnosis of Idiopathic Short Stature (ISS). Since this is no longer a covered benefit, this will be a benefit denial.

#### **RATIONALE:**

1. The technology must have final approval from appropriate government regulatory bodies (e.g., the FDA). The FDA has approved the labeled use of human growth hormone for specific indications.
2. The scientific evidence must permit conclusions concerning the effect of the technology on health outcomes. Numerous clinical trials have been published to demonstrate the efficacy and safety of GH for specific conditions (i.e., burn patients). The evidence is insufficient to permit conclusions concerning the effect of GH therapy on health outcomes on geriatric patients, patients with non-GH deficient short stature, and patients with cardiac disease, critically- ill patients, or other conditions where anabolic therapy has been suggested to counteract acute illness.

## Pharmacy Management Drug Policy

### Growth Hormone

3. The technology must improve net health outcomes; and
4. The technology must be as beneficial as any established alternatives.
5. Published clinical trials have demonstrated that GH stimulates growth in children, increases body weight, increases lean body mass, decreases fat mass, increases bone density and stimulates bone turnover when used in adults for specific conditions.
6. The improvement must be attainable outside the investigational settings. GH therapy has been proven to improve net health outcomes outside the investigational setting in specific instances (i.e. children and adults with GHD, AIDS wasting).

#### **UPDATES:**

<b>Date:</b>	<b>Revision:</b>
<b>3/19</b>	<b>Revised</b>
2/19	P&T Approval
9/18	Reviewed
11/17	Reviewed
12/16	Reviewed
12/15	Reviewed
12/14	Revised
8/14	Reviewed
8/13	Revised/P&T Approval
4/10	Reviewed
6/09	Reviewed
9/08	Revised
7/09	Revised

#### **REFERENCES:**

1. American Academy of Pediatrics Committee on Drugs and Committee on Bioethics, "Considerations Related to the Use of Recombinant Human Growth Hormone in Children", Pediatrics, January 1997, 99(1): 122 – 9.
2. Blue Cross Blue Shield Association, "Human Growth Hormone", Medical Policy Reference Manual, #5.01.06, Issued 1997.
3. Carroll PV, et al, "Growth Hormone Deficiency in Adulthood and the Effects of Growth Hormone Replacement: A Review", Journal of Clinical Endocrinology and Metabolism, 1998, 83(2): 382-94.
4. Ellegard L, et al, "Low Dose Recombinant Human Growth Hormone Increases Body Weight and Lean Mass in patients with Short Bowel Syndrome", Annals of Surgery, January 1997, 225(1): 88-96.
5. Jeschke MG, et al, "Recombinant Human Growth Hormone Treatment in Pediatric Burn Patients and its Role During the Hepatic Acute Phase Response", Critical Care Medicine, May 28, 2000, (5): 1578-84.
6. Osterziel KJ, et al, "Randomized, Double-Blinded, Placebo, Controlled Trial of Human Growth Hormone in Patients with Chronic Heart failure Due to Dilated Cardiomyopathy", Lancet, April 25, 1998, 351(9111): 1233-7.



## Pharmacy Management Drug Policy

### Growth Hormone

7. Scolapio JS, et al, "Effect of Growth Hormone, Glutamine and Diet on Adaptation in Short Bowel Syndrome: A Randomized, Controlled Study", Gastroenterology, October 1997, 113(4): 1074-81.
8. Ramirez RJ, et al, "Growth Hormone Treatment in Pediatric Burns: A Safe Therapeutic Approach", Annals of Surgery, October 1998, 228(4): 439-48.
9. Slonim AE, et al, "A Preliminary Study of Growth Hormone Therapy for Crohn's Disease", NEJM, June 1, 2000, 342(22): 1633-7.
10. Takala J, et al, "Increased Mortality Associated with Growth Hormone treatment in Critically Ill Adults", NEJM, September 9, 1999, 341(11): 785-92.
11. The Endocrine Society, "Invited Report of a Workshop. Consensus Guidelines for the Diagnosis and Treatment of Adults with Growth Hormone Deficiency: Summary Statement of the Growth Hormone Research Society", Journal of Clinical Endocrinology and Metabolism, 1998, 83(2): 379-81.
12. The Endocrine Society, "Consensus Guidelines for the Diagnosis and Treatment of Growth Hormone Deficiency in Childhood and Adolescence: Summary Statement of the GH Research Society", Journal of Clinical Endocrinology and Metabolism, 2000, 85(11): 3990-93.
13. Badaru A, et al, "Alternatives to growth hormone stimulation testing in children", Trends in Endocrinology and Metabolism, Aug 2004, Vol15 No. 6: 252-258
14. Cianfiarani S, et al, "Height velocity and IGF-1 assessment in the diagnosis of childhood onset GH insufficiency: do we still need a second GH stimulation test?", Clinical Endocrinology 2002, 57: 161-167
15. Zorbtive package insert, EMD Serono, Inc. Revised January 2012. – Accessed online December 2016
16. Genotropin package insert, Pfizer Inc. Revised May 2015 – Accessed online November 2017
17. Humatrope package insert , Eli Lilly & Company, Revised July 30, 2014 - Accessed online November 2017
18. Kojima, M et al. Ghrelin: structure and function. Physiology Rev. 2005 April. 85(2), 495-522.
19. Sas T, Dewaal W, Mulder P, et al. Growth hormone treatment in children with short stature born small for gestational age: 5-year results of a randomized, double-blind, dose-response trial. J Clin Endocrinol Metab 1999; 84;3064-3070
20. Rosilio M, Carel JC, Ecosse E, et al. Adult height of prepubertal children born small for gestational age treated with GH Eur J Endocrinol 2005;152:835-843
21. Corneli g, Gasco V, Prodam F, et al. Growth hormone levels in the diagnosis of growth hormone deficiency in adulthood Pituitary 2007 Jun; 10(2):141-149
22. Reiter EO A brief review of the addition of gonadotropin releasing hormone agonists (GnRH-Ag) to growth hormone (GH) treatment of children with idiopathic growth hormone deficiency: Previously published studies from America. Mol Cell Endocrinol 2006 Jul;254-255:221-5
23. Wilson TA, Rose SR, Cohen P, et al. Update of guidelines for the use of growth hormone in children: the Lawson Wilkins Pediatric Endocrinology Society Drug and Therapeutics Committee. J Pediatr. 2003;143:415-421

## Pharmacy Management Drug Policy

### Growth Hormone

24. Molitch ME, Clemmons DR, Malozowski S, et al; Endocrine Society's Clinical Guidelines Subcommittee; Stephens PA. Evaluation and treatment of adult growth hormone deficiency: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab.* 2006;91:1621-1634. Available at: <http://www.endo-society.org/publications/guidelines/index.cfm> Accessed 08/2008.
25. Blum WF, Crowe BJ, Quigley CA, et al; Rappold For The Shox Study Group G. Growth hormone is effective in treatment of short stature associated with short stature homeobox-containing gene: two-year results of a randomized, controlled, multi-center trial. *J Clin Endocrinol Metab.* 2007;92:219-228
28. American Association of Clinical Endocrinologists Position Statement Growth Hormone Usage in Short Children December 2003
28. Omnitrope full prescribing information. Sandoz Inc. Revised October 2014 - Accessed online November 2017
29. T. Romer et al. Seven Years of Safety and Efficacy of the Recombinant Human Growth Hormone Omnitrope in the Treatment of Growth Hormone Deficient Children: Results of a Phase III Study. *Horm Res* 2009;72:359–369.
30. Cook DM et al. American Association Of Clinical Endocrinologists Medical Guidelines For Clinical Practice For Growth Hormone Use In Growth Hormone-Deficient Adults And Transition Patients – 2009 update. *Endocrine Practice.* Vol 15 (Suppl 2) September/October 2009
31. Wilson TA et al. Update Of Guidelines For The Use Of Growth Hormone In Children: The Lawson Wilkins Pediatric Endocrinology Society Drug And Therapeutics Committee. *The Journal of Pediatrics.* October 2003;143(4);415-21.
32. US Food and Drug Administration: FDA Drug Safety Communication: Safety review update of Recombinant Human Growth Hormone (somatropin) and possible increased risk of death <http://www.fda.gov/Drugs/DrugSafety/ucm265865.htm>; update 8/4/2011
33. Bell J, Parker KL, Swinford RD, Hoffman AR, Maneatis T, Lippe B. Long-term safety of recombinant human growth hormone in children. *J Clin Endocrinol Metab.* 2010;95:167-77.
34. Up to Date: Growth Hormone Treatment for Idiopathic Short Stature; accessed online at [uptodate.com](http://uptodate.com); updated March 2016 - Accessed online December 2016
35. Schambelan M, Mulligan K, Grunfeld C, Daar ES, LaMarca A, Kotler DP, Wang J, Bozzette SA, Breitmeyer JB “Recombinant human growth hormone in patients with HIV-associated wasting. A randomized, placebo-controlled trial. Serostim Study Group”. *Ann Intern Med.* 1996;125(11):873
36. Pediatrics Vol. 114 No. Supplement 6 November 1, 2004; pp. 1478 -1482 (doi:10.1542/peds.2004-1721P)