

# MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	LIMB PNEUMATIC COMPRESSION DEVICES FOR VENOUS THROMBOEMBOLISM PROPHYLAXIS
Policy Number	1.01.51
Category	Technology Assessment
Effective Date	06/27/13
Revised Date	06/26/14, 10/15/15, 06/16/16, 07/20/17, 07/19/18, 07/18/19
Product Disclaimer	<ul style="list-style-type: none"> <li>• If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</li> <li>• If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit.</li> <li>• If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</li> </ul>

## POLICY STATEMENT

- I. Based on our criteria, pneumatic compression devices have been medically proven to be effective and are considered **medically appropriate** when used in the home for deep vein thrombosis (DVT) prophylaxis after major surgery, including major orthopedic surgery, in patients when pharmacological prophylaxis is contraindicated. Use of pneumatic compression device is allowed for up to 30 days post-operatively.
- II. Based upon our criteria and assessment of peer-reviewed literature, use of pneumatic compression devices for prevention of deep vein thrombosis, other than as described in Policy Statement I, is considered **not medically necessary**.

*Refer to Corporate Medical Policy 1.01.17 Pneumatic Compression Devices/Lymphedema Pumps*

## POLICY GUIDELINES

- I. Major orthopedic surgery includes total hip arthroplasty (THA), total knee arthroplasty (TKA), or hip fracture surgery (HFS).
- II. American College of Chest Physicians (ACCP) guidelines on prevention of VTE in orthopedic surgery patients list the following general risk factors for bleeding:
  - A. Previous major bleeding (and previous bleeding risk similar to current risk),
  - B. Severe renal failure,
  - C. Concomitant antiplatelet agent, and
  - D. Surgical factors: history of or difficult-to-control surgical bleeding during the current operative procedure, extensive surgical dissection, and revision surgery.
- III. The Federal Employee Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

## DESCRIPTION

Pneumatic compression devices have been utilized in the post-operative period for prevention of deep vein thrombosis. The devices include various types of wraps for the arms or legs and a programmable control module. The wraps of some of these devices are capable of providing cooling or heating to the extremity. Examples of devices which can be used in

## **Medical Policy: LIMB PNEUMATIC COMPRESSION DEVICES FOR VENOUS THROMBOEMBOLISM PROPHYLAXIS**

**Policy Number: 1.01.51**

**Page: 2 of 5**

the home after discharge include the VascuTherm2 (Thermotek, Inc) and the Triple Play VT® (Compression Solutions, Inc).

### **RATIONALE**

Anticoagulation is the mainstay of DVT prophylaxis after major surgery and is sometimes continued into the outpatient setting. Treatment with pneumatic compression devices may offer an additional benefit when used in conjunction with anticoagulation in the inpatient setting, but is not commonly used in the outpatient setting. Studies involving the use of compression devices post-operatively are limited; the studies are often small, nonrandomized with considerable variation in the comparison studies by type of compression stocking and intermittent compression device used, patient group, DVT detection method, and prophylaxis protocol. Many of the studies are in the setting of the hospital rather than outpatient, consequently conclusions from the hospital setting may not be able to be applied to the outpatient setting. This is due to differences in the levels of mobility in the hospital versus the outpatient setting, which may change the risk for DVT. Also, the use of pneumatic compression devices in the hospital can be more highly controlled and monitored. In the outpatient setting, there are questions about the degree of compliance with the devices, including the ability to correctly use them in the absence of professional supervision.

In 2011, the American Academy of Orthopaedic Surgeons (AAOS) published an updated guideline on prevention of venous thromboembolism in patients undergoing elective hip and knee arthroplasty. The work group recommended the use of pharmacologic agents and/or mechanical compressive devices for the prevention of venous thromboembolism in patients undergoing elective hip or knee arthroplasty who are not at elevated risk beyond that of the surgery itself for venous thromboembolism or bleeding (moderate recommendation), and who have also had a previous venous thromboembolism (consensus recommendation). For those patients undergoing elective hip or knee arthroplasty, who also have a known bleeding disorder (e.g., hemophilia) and/or active liver disease, mechanical compressive devices for preventing venous thromboembolism is recommended (consensus recommendation).

The American College of Chest Physicians (ACCP) (2012) recommended the use of limb compression devices in non-orthopedic general and abdominal-pelvic surgical patients, in patients who were classified as low risk for VTE and moderate risk for VTE and high-risk for bleeding rather than no prophylaxis and as an option for those individuals with moderate and high risk for VTE and who were not at high risk for bleeding in addition to pharmacologic prophylaxis.

In 2007, the American College of Obstetricians-Gynecologists (ACOG) published a practice bulletin on prevention of DVT and PE after gynecologic surgery. As with the ACCP recommendations, described above, prophylaxis recommendations varied according to patient risk level. For patients at moderate and high risk of DVT, intermittent pneumatic compression was one of the recommended options for DVT prophylaxis. For patients at highest risk i.e., older than 60 years plus prior VTE, cancer or molecular hypercoagulable state, IPC or graduated compression stockings plus LDUH or LMWH was recommended as a prophylaxis option. The devices should be used continuously until ambulation and discontinued only at the time of hospital discharge except for the highest risk patients, who should consider prophylaxis for two-four weeks after discharge.

A Cochrane review by Naccarato, et al. (2010), identified two clinical trials and two small studies using gradient compression stockings (GCS) or intermittent pneumatic compression (IPC) devices for DVT prophylaxis in stroke patients. The authors concluded that evidence from randomized trials does not support the routine use of GCS to reduce the risk of DVT after acute stroke. There is insufficient evidence to support the routine use of IPC to reduce the risk of DVT in acute stroke and further larger randomized studies of IPC are needed to reliably assess the balance of risks and benefits of this intervention.

### **CODES**

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- ***CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.***
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*

**Medical Policy: LIMB PNEUMATIC COMPRESSION DEVICES FOR VENOUS THROMBOEMBOLISM  
PROPHYLAXIS**

**Policy Number: 1.01.51**

**Page: 3 of 5**

**CPT Codes**

<b>Code</b>	<b>Description</b>
No Codes	

*Copyright © 2019 American Medical Association, Chicago, IL*

**HCPCS Codes**

<b>Code</b>	<b>Description</b>
E0650	Pneumatic Compressor, nonsegmental home model
E0651	Pneumatic compressor, segmental home model without calibrated gradient pressure
E0652	Pneumatic Compressor, segmental home model with calibrated gradient pressure
E0660	Nonsegmental pneumatic appliance for use with pneumatic compressor, full leg
E0666	half leg
E0667	Segmental pneumatic appliance for use with pneumatic compressor, full leg
E0669	half leg
E0671	Segmental gradient pressure pneumatic appliance, full leg
E0673	half leg
E0675	Pneumatic compression device, high pressure, rapid inflation/deflation cycle for arterial insufficiency (unilateral or bilateral system)
E0676	Intermittent limb compression device (includes all accessories), not otherwise specified

**ICD10 Codes**

<b>Code</b>	<b>Description</b>
M05.00-M05.09	Felty's syndrome (code range)
M05.20-M05.29	Rheumatoid vasculitis with rheumatoid arthritis (code range)
M05.30-M05.39	Rheumatoid heart disease with rheumatoid arthritis (code range)
M05.40-M05.49	Rheumatoid myopathy with rheumatoid arthritis (code range)
M05.50- M05.579	Rheumatoid polyneuropathy with rheumatoid arthritis (code range)
M05.60-M05.79	Rheumatoid arthritis of unspecified site with or without involvement of other organs and systems (code range)
M05.80-M06.09	Rheumatoid arthritis with or without rheumatoid factor (code range)
M06.1	Adult-onset Still's disease
M06.2.0-M06.29	Rheumatoid bursitis (code range)
M06.30-M06.39	Rheumatoid nodule (code range)
M06.4	Inflammatory polyarthropathy
M06.80-M06.9	Other specified rheumatoid arthritis (code range)
M08.00-M08.29	Unspecified juvenile rheumatoid arthritis (code range)
M08.3	Juvenile rheumatoid polyarthritis (seronegative)
M08.40-M08.48	Pauciarticular juvenile rheumatoid arthritis (code range)
M08.80-M08.99	Other juvenile arthritis (code range)
M12.00- M12.569	Chronic post-rheumatic arthropathy (Jaccoud) (code range)
M15.0-M19.93	Osteoarthritis (code range)
M80.051A- M80.859A	Osteoporosis (code range)
M84.451A- M84.659A	Pathological fracture, hip or femur, initial encounter for fracture (code range)

## Medical Policy: LIMB PNEUMATIC COMPRESSION DEVICES FOR VENOUS THROMBOEMBOLISM PROPHYLAXIS

Policy Number: 1.01.51

Page: 4 of 5

Code	Description
S72.001A- S79.099A	Fracture of lower extremity, initial encounter for fracture type (code range)
Z47.1	Aftercare following joint replacement surgery
Z96.641- Z96.659	Presence of artificial lower extremity joint (code range)

### REFERENCES

\*Akpınar EE, et al. Does thromboprophylaxis prevent venous thromboembolism after major orthopedic surgery? J Bras Pneumol 2013 May-Jun;39(3):280-6.

Afshari A, et al. European guidelines on perioperative venous thromboembolism prophylaxis: Mechanical prophylaxis. Eur J Anaesthesiol. Feb 2018;35(2):112-115.

Ahmed AB, et al. European guidelines on perioperative venous thromboembolism prophylaxis: Cardiovascular and thoracic surgery. Eur J Anaesthesiol. Feb 2018;35(2):84-89

BlueCross BlueShield Association. Lymphedema pumps. Medical Policy Reference Manual Policy #1.01.18.2019 Mar 14.

BlueCross BlueShield Association. Postsurgical home use of limb compression devices for venous thromboembolism prophylaxis. Medical Policy Reference Manual Policy #1.01.28. 2019 Mar 14.

\*Colwell CW, et al. A mobile compression device for thrombosis prevention in hip and knee arthroplasty. J Bone Joint Surg Am 2014 Feb 5;96(3):177-83.

\*Committee on Practice Bulletins-Gynecology, American College of Obstetricians and Gynecologists. ACOG Practice Bulletin No. 84: Prevention of deep vein thrombosis and pulmonary embolism. Obstet Gynecol 2007;110(2 Pt 1):429-40.

Domeij-Arverud E, et al. Intermittent pneumatic compression reduces the risk of deep vein thrombosis during post-operative lower limb immobilization: a prospective randomized trial of acute ruptures of the Achilles tendon. Bone Joint J 2015 May;97-B(5):675-80.

Falck-Ytter Y, et al. Prevention of VTE in orthopedic surgery patients: antithrombotic therapy and prevention of thrombosis. 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012;141:e278S-e325S.

Faraoni D, et al. European guidelines on perioperative venous thromboembolism prophylaxis: Neurosurgery. Eur J Anaesthesiol. Feb 2018;35(2):90-95.

\*Feist WR, et al. Problems with measuring compression device performance in preventing deep vein thrombosis. Throm Res 2011 Sep;128(3):207-9.

Feng JP, et al. Efficacy of intermittent pneumatic compression for venous thromboembolism prophylaxis in patients undergoing gynecologic surgery: a systematic review and meta-analysis. Oncotarget 2017;8(12):20371-79.

Flierl MA, et al. Venous thromboembolism prophylaxis after total joint arthroplasty. Orthopedics 2015 Apr 1;38(4):252-63.

Gould MK, et al. Prevention of VTE in nonorthopedic surgical patients: Antithrombotic Therapy and Prevention of Thrombosis. 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012;141(2 Suppl):e227S-77S.

## **Medical Policy: LIMB PNEUMATIC COMPRESSION DEVICES FOR VENOUS THROMBOEMBOLISM PROPHYLAXIS**

**Policy Number: 1.01.51**

**Page: 5 of 5**

\*Guyatt GH, et al. American College of Chest Physicians Antithrombotic therapy and Prevention of Thrombosis Panel. Executive summary: Antithrombotic Therapy and Prevention of Thrombosis, 9<sup>th</sup> ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest 2012;141(2 Suppl):7S-47S.

Kakkos SK, et al. Combined intermittent pneumatic leg compression and pharmacological prophylaxis for prevention of venous thromboembolism. Cochrane Database Syst Rev. Sep 07 2016;9:CD005258.

O'Connell S, et al. The use of intermittent pneumatic compression in orthopedic and neurosurgical postoperative patients: a systematic review and meta-analysis. Ann Surg 2016 May;263(5):888-98.

Sadaghianloo N, et al. The efficacy of intermittent pneumatic compression in the prevention of lower extremity deep venous thrombosis. J Vasc Surg Venous Lymphat Disord 2016 Apr;4(2):248-56.

Saunders R, et al. Intermittent pneumatic compression is a cost-effective method or orthopedic postsurgical venous thromboembolism prophylaxis. Clinicoecon Outcomes Res. 2018 Apr 19;10:231-241.

Takahashi H, et al. Venous thromboembolism after elective shoulder surgery: a prospective cohort study of 175 patients. J Shoulder Elbow Surg 2014 May;23(5):605-12.

Torrejon TR, et al. A comparative cost-effectiveness analysis of mechanical and pharmacological VTE prophylaxis after lower limb arthroplasty in Australia. J Orthop Surg Res. 2019 Apr 2;14(1):93.

Zareba P, et al. Meta-analysis of randomized trials comparing combined compression and anticoagulation with either modality alone for prevention of venous thromboembolism after surgery. Br J Surg 2014 Aug;101(9):1053-62.

\*Key Article

### **KEY WORDS**

Venodyne, VascuTherm2, Triple Play VT®

### **CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS**

There is currently a National Coverage Determination (NCD) and a Local Coverage Determination (LCD) for Pneumatic Compression Devices. Please refer to the following websites for Medicare Members:

NCD SITE: <http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=225&ncdver=1&bc=AgAAgAAAAAAA&>

LCD SITE: [https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33829&ver=38&CntrctrSelected=137\\*1&Cntrctr=137&s=41&DocType=Active&bc=AggAAAIIBA AAA&](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33829&ver=38&CntrctrSelected=137*1&Cntrctr=137&s=41&DocType=Active&bc=AggAAAIIBA AAA&)