

MEDICAL POLICY



MEDICAL POLICY DETAILS	
Medical Policy Title	PELVIC FLOOR ELECTRICAL STIMULATION AS A TREATMENT FOR URINARY OR FECAL INCONTINENCE
Policy Number	1.01.19
Category	Technology Assessment
Effective Date	09/16/99
Revised Date	08/16/01, 07/18/02, 05/21/03, 06/17/04, 06/16/05, 06/15/06, 05/17/07, 05/14/08, 09/18/08, 09/17/09, 09/16/10, 09/15/11
Archived Date	09/20/12
Edited Date	09/19/13, 09/18/14, 09/17/15, 07/21/16, 07/20/17, 07/19/18, 07/18/19
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

Based upon our criteria and assessment of peer-reviewed literature, there is no evidence that pelvic floor electrical stimulation demonstrates improvement in patient outcomes. Therefore, pelvic floor electrical stimulation (PFES) for the treatment of urinary or fecal incontinence is considered **not medically necessary**.

Refer to Corporate Medical Policy #1.01.01 regarding Electrical Stimulation - Transcutaneous Electrical Nerve (TENS), H-Wave and Interferential Stimulators.

Refer to Corporate Medical Policy #7.01.10 Sacral Nerve Stimulation for Pelvic Floor Dysfunction.

POLICY GUIDELINES

The Federal Employee Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

DESCRIPTION

Urinary incontinence is defined by the International Continence Society (ICS) as “a condition in which involuntary loss of urine is a social or hygienic problem.” The National Institute of Health (NIH) statistics indicate urinary incontinence is estimated to affect 10-12 million people in the United States, two thirds of who are female.

Pelvic floor electrical stimulation (PFES) has been advocated as a treatment of urinary stress incontinence, urge incontinence and incontinence due to detrusor instability. PFES is also being investigated as a treatment modality for patients with fecal incontinence due to pelvic floor dysfunction and has also been proposed as a non-invasive alternative to surgical intervention for patients with damage to the anal sphincter.

PFES is the application of electrical current to the pudendal nerve. This electrical stimulation causes reflex contraction of the pelvic floor musculature (detrusor/bladder muscle and levator ani muscle). PFES is applied to the body using skin electrodes around the anus or by vaginal or rectal sensors (probes). PFES may be used alone or in conjunction with biofeedback or pelvic floor muscle exercises. The goal of PFES is to regain volitional control of the pelvic floor muscles through their passive activation. The patient should progress to regaining voluntary control of muscle contraction without stimulation, increasing the strength of pelvic floor muscles thus, and eliminating urinary leakage.

Medical Policy: PELVIC FLOOR ELECTRICAL STIMULATION AS A TREATMENT FOR URINARY OR FECAL INCONTINENCE

Policy Number: 1.01.19

Page: 2 of 3

RATIONALE

While case series have reported promising outcomes of pelvic floor electrical stimulation, the evidence from these case series tends to overestimate the treatment effect. These studies do not account for placebo effects or for dropouts. Many of the studies investigating electrical stimulation as a treatment of urinary or fecal incontinence combined biofeedback and/or pelvic floor muscle exercises with electrical stimulation as the intervention, which makes it difficult to determine the independent effect of electrical stimulation alone. Published randomized, controlled clinical trials investigating this treatment modality have reported inconsistent and/or inconclusive results. The evidence is insufficient to determine the effectiveness of pelvic floor electrical stimulation on urinary incontinence and fecal incontinence. Most of the published studies have not measured the effect of these devices on pelvic muscle strength, which is a measurement of the effectiveness of pelvic floor stimulation. There is insufficient evidence from clinical trials to determine whether electrical stimulation is more effective than pelvic floor muscle exercises or even sham electrical stimulation.

CODES

- Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.
- Code Key: Experimental/Investigational = (E/I), Not medically necessary/ appropriate = (NMN).

CPT Codes

Code	Description
97014*	Physical medicine and rehabilitation - application of a modality to one or more areas; electrical stimulation (unattended) (*NMN for N39.3, N39.41-N39.498, R15.0-R15.9, R32)
97032*	application of a modality to one or more areas; electrical stimulation (manual), each 15 minutes (*NMN for N39.3, N39.41-N39.498, R15.0-R15.9, R32)

Copyright © 2019 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
E0740 (NMN)	Non-implanted pelvic floor electrical stimulator, complete system

ICD10 Codes

Code	Description
N39.3	Stress incontinence (female) (male)
N39.41-N39.498	Other specified urinary incontinence (code range)
R15.0-R15.9	Fecal incontinence (code range)
R32	Unspecified urinary incontinence

REFERENCES

Andersen CA, et al. Conservative management for postprostatectomy urinary incontinence. Cochrane Database Syst Rev 2015 Jan 20;1:CD001843.

BlueCross BlueShield Association. Pelvic Floor Stimulation as a Treatment of Urinary Incontinence. Medical Policy Reference Manual Policy #1.01.17. 2018 Aug 09.

Medical Policy: PELVIC FLOOR ELECTRICAL STIMULATION AS A TREATMENT FOR URINARY OR FECAL INCONTINENCE

Policy Number: 1.01.19

Page: 3 of 3

- *BlueCross BlueShield Association. Technology Evaluation Center (TEC). Pelvic floor electrical stimulation in the treatment of urinary incontinence in adults. 2000 Jun;15(2).
- *Castro RA, et al. Single-blind, randomized, control trial of pelvic floor muscle training, electrical stimulation, vaginal cones, and no active treatment in the management of stress urinary incontinence. Clinics 2008;64:465-72.
- *Gilling PJ, et al. A double-blind randomized controlled trial of electromagnetic stimulation of the pelvic floor vs sham therapy in treatment of women with stress urinary incontinence. BJU Int 2009;103(10):1386-90.
- *Goode PS, et al. Behavioral therapy with or without biofeedback and pelvic floor electrical stimulation for persistent postprostatectomy incontinence: a randomized controlled trial. JAMA 2011 Jan 12;305(2):151-9.
- *Goode PS, et al. Effect of behavioral training with or without pelvic floor electrical stimulation on stress incontinence in women. JAMA 2003 Jul;290(3):345-52.
- *Mahoney RT, et al. Randomized clinical trial of intra-anal electromyographic biofeedback physiotherapy with intra-anal electromyographic biofeedback augmented with electrical stimulation of the anal sphincter in early postpartum fecal incontinence. Am J Obstet Gynecol 2004 Sep;191(3):885-90.
- National Institute for Health and Care Excellence (NICE). Urinary incontinence and pelvic organ prolapse in women: management. NG123. 2019 Apr [<https://www.nice.org.uk/guidance/ng123>] accessed 6/10/19.
- *Osterberg A, et al. Randomized clinical trial comparing conservative and surgical treatment of neurogenic faecal incontinence. Br J Surg 2004 Sep;91(9):1131-7.
- Shamliyan TA, et al. Nonsurgical treatments for urinary incontinence in adult women: diagnosis and comparative effectiveness. Comparative Effectiveness Reviews, No. 36. 2012 Apr [<https://effectivehealthcare.ahrq.gov/topics/urinary-incontinence-treatment/research>] accessed 6/10/19.
- *Spruijt J, et al. Vaginal electrical stimulation of the pelvic floor: a randomized feasibility study in urinary incontinent elderly women. ACTA Obstet Gynecol Scand 2003 Nov;82(11):1043-8.
- Stewart F, et al. Electrical stimulation with non-implanted devices for stress urinary incontinence in women. Cochrane Database Syst Rev 2017 Dec 22;12:CD012390.
- *Vonthein R, et al. Electrical stimulation and biofeedback for the treatment of fecal incontinence: a systematic review. Int J Colorectal Dis 2013; 28(11):1567-77.
- *Wille S, et al. Pelvic floor exercises, electrical stimulation and biofeedback after radical prostatectomy: results of a prospective randomized trial. J Urol 2003 Aug;170(2 Pt 1):490-3.
- *Zhu YP, et al. Pelvic floor electrical stimulation for postprostatectomy urinary incontinence: a meta-analysis. Urology 2012 Mar;79(3):552-5.

*Key Article

KEY WORDS

Intravaginal electrical stimulation.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently a National Coverage Determination (NCD) for Non-Implantable Pelvic Floor Electrical Stimulator. Please refer to the following NCD website for Medicare Members: <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=231&ncdver=2&bc=AgAAgAAAAAA&>.