

# MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	<b>POSITIVE AIRWAY PRESSURE DEVICES: CPAP, BiPAP, APAP AND NONINVASIVE POSITIVE PRESSURE VENTILATORS</b>
Policy Number	<b>1.01.06</b>
Category	<b>Equipment/Supplies</b>
Effective Date	<b>10/18/01</b>
Revised Date	<b>01/17/02, 02/20/03, 12/18/03, 01/20/05, 04/21/05, 03/16/06, 04/26/07, 06/26/08, 02/26/09, 04/28/11, 04/26/12, 04/27/17, 08/25/17, 04/26/18, 6/27/19</b>
Archived Date	<b>04/25/13 - 4/28/16</b>
Edited Date	<b>04/24/14, 06/26/14, 08/28/14, 04/23/15, 04/28/16</b>
Product Disclaimer	<ul style="list-style-type: none"> <li>• <i>If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</i></li> <li>• <i>If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit.</i></li> <li>• <i>If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</i></li> </ul>

## POLICY STATEMENT

### I. Obstructive Sleep Apnea in adults:

Based upon our criteria and assessment of the peer-reviewed literature, *Continuous positive airway pressure (CPAP)* (HCPCS: E0601) has been medically proven to be effective and therefore **medically appropriate** for the treatment of obstructive sleep apnea *for an initial 90 day trial period* in adults who meet the following criteria:

- A. Polysomnography (PSG) or home sleep testing (HST) results documenting an apnea-hypopnea index (AHI) or respiratory disturbance index (RDI) of greater than 5 and less than 14 respiratory events per hour with a minimum of 10 respiratory events and associated symptoms of obstructive sleep apnea (OSA) (e.g., excessive daytime sleepiness, impaired cognition, and insomnia) or documented cardiovascular diseases, including hypertension, ischemic heart disease, or history of stroke;  
OR
- B. Polysomnography or HST results documenting an AHI or RDI of greater than 15 with a minimum of 30 respiratory events;  
AND
- C. A compliance support plan between the treating physician and DME supplier has been established.
- D. Continuation of coverage after 90 day trial period when the following has been demonstrated:
  1. Improved AHI and symptom resolution during trial period); and
  2. Compliance has been demonstrated as defined by the use of the device for 70% of the nights for an average of 4 or more hours per 24 hour period during a consecutive 30 day period.
- E. Patients who do not meet the initial 90 day trial for continuation of coverage are eligible to re-qualify for a CPAP device but must have a face-to-face clinical re-evaluation by the treating physician to determine the etiology of the failure to respond to CPAP therapy (e.g., documentation of failure of symptoms to resolve or improper fit of device and re-education of the patient regarding proper use of the equipment or re-fitting of masks).
- F. *Bilevel Positive Airway Pressure (BPAP)* (HCPCS: E0470 or E0471) with or without rate control is considered a medically appropriate treatment option *for an initial 90 day trial period* in adults diagnosed with obstructive sleep apnea who have failed CPAP or not tolerated CPAP and there is documentation of demonstration of resolution/control of respiratory events with improved tolerance using BPAP.
- G. Continuation of coverage after 90 day trial period when the following has been demonstrated:
  1. Improved AHI and symptom resolution during trial period); and

**Medical Policy: POSITIVE AIRWAY PRESSURE DEVICES: CPAP, BiPAP, APAP AND NONINVASIVE POSITIVE PRESSURE VENTILATORS**

**Policy Number: 1.01.06**

**Page: 2 of 10**

2. Compliance has been demonstrated as defined by the use of the device for 70% of the nights for an average of 4 or more hours per 24 hour period during a consecutive 30 day period.

**II. Obstructive Sleep Apnea in children:**

Based upon our criteria and assessment of peer-reviewed literature, CPAP has been medically proven to be effective and therefore **medically appropriate** for the treatment of obstructive sleep apnea in children for the following indications:

- A. Failure of adenotonsillectomy to relieve OSA symptoms or a contraindication to surgical intervention or for whom there is a strong preference for a nonsurgical approach; AND
- B. Polysomnography results documenting an AHI or RDI of 5 or greater and associated symptoms of OSA. Common OSA symptoms in children include (but are not limited to) habitual snoring, disturbed sleep, and daytime neurobehavioral problems such as hyperactivity. Daytime sleepiness may be present, but is less common in children. Children with an AHI or RDI greater than 2 but less than 5 may be considered for CPAP if significant daytime symptoms exist; OR
- C. Polysomnography results demonstrating an AHI or RDI within normal ranges, but the child exhibits episodes of hypercarbia based on end-tidal CO<sub>2</sub> measurements greater than 53 mm Hg or the end-tidal CO<sub>2</sub> is greater than 50 mmHg for 10 – 24% of the sleep time. The child must also exhibit associated symptoms of OSA.
- D. The diagnostic criteria for pediatric obstructive sleep apnea (OSA) as defined by the American Academy of Sleep Medicine (AASM) and International Classification of Sleep Disorders (ICSD) is below. *All* of the following criteria should be present for a child to be diagnosed with OSA:
  1. Snoring, labored breathing, or obstructed breathing during the child's sleep.
  2. One or more of the following: paradoxical inward rib cage motion during inspiration, movement arousals, diaphoresis, neck hyperextension during sleep, excessive daytime sleepiness, hyperactivity, aggressive behavior, slow growth, morning headaches, or secondary enuresis.
  3. Polysomnography reveals one or more obstructive apneas or hypopneas per hour of sleep (e.g., an apnea hypopnea index greater than 1 event per hour).
  4. Polysomnography demonstrates either of the following:
    - a. frequent arousals from sleep associated with increased respiratory effort, oxyhemoglobin desaturation associated with apnea, hypercapnia during sleep, or markedly negative esophageal pressure swings; or
    - b. periods of hypercapnia, oxyhemoglobin desaturation, or both during sleep that are associated with snoring, paradoxical inward rib cage motion during inspiration, and either frequent arousals from sleep or markedly negative esophageal pressure swing.

**III. Snoring:**

Based upon our criteria and the lack of peer-reviewed literature, CPAP is **not medically necessary** for the treatment of snoring without accompanying OSA.

**IV. Central Sleep or Complex Sleep Apnea:**

- A. *Bilevel Positive Airway Pressure (BPAP)* (HCPCS: E0470 or E0471) with or without rate control is considered a **medically appropriate** treatment option for *an initial 90 day trial period* in adults diagnosed with central sleep or complex sleep apnea who have failed CPAP or not tolerated CPAP and who meet the following criteria :
  1. central hypopnea/apnea greater than 50% of the total apnea hypopnea rate; or
  2. central hypopnea/apnea rate/index greater than 5 events per hour; or
  3. significant improvement of oxygenation while breathing the patient's prescribed FiO<sub>2</sub>; or
  4. documentation of demonstration of resolution/control of respiratory events with improved tolerance using BPAP.
- B. Continuation of coverage after 90 day trial period when the following has been demonstrated:
  1. Improved AHI and symptom resolution during trial period); and
  2. Compliance has been demonstrated as defined by the use of the device for 70% of the nights for an average of 4 or more hours per 24 hour period during a consecutive 30 day period.

**Medical Policy: POSITIVE AIRWAY PRESSURE DEVICES: CPAP, BiPAP, APAP AND NONINVASIVE POSITIVE PRESSURE VENTILATORS**

**Policy Number: 1.01.06**

**Page: 3 of 10**

V. Chronic respiratory failure:

Based upon our criteria and assessment of the peer-reviewed literature, *BPAP*, when used as noninvasive ventilatory support with or without the back-up rate feature, has been medically proven to be effective and therefore **medically appropriate** for an initial 90 day trial period in adults who have failed CPAP and who meet the following indications:

- A. Restrictive thoracic disorders (e.g. neuromuscular disease, thoracic cage abnormalities) with documented:
  - 1. oxygen desaturations less than or equal to 88% for at least 5 continuous minutes during sleep oximetry (minimum recording time of 2 hours) while the patient is either breathing room air or prescribed FIO<sub>2</sub> as applicable, or
  - 2. arterial blood gas PaCO<sub>2</sub> greater than or equal to 45 mm Hg while the patient is awake and breathing either room air or prescribed FIO<sub>2</sub> as applicable; or
  - 3. a documented decrease in forced vital capacity (FVC) or vital capacity (VC) to 50% of predicted; or
  - 4. a maximal inspiratory pressure of less than 60 cm H<sub>2</sub>O; or
  - 5. there is a symptomatic respiratory disease impairing activities of daily living;AND
- 6. COPD does not contribute to the patient's pulmonary limitation.
- B. Severe chronic pulmonary disease with:
  - 1. arterial blood gas PaCO<sub>2</sub> is greater than or equal to 52 mm Hg while the patient is awake and is breathing either breathing room air or prescribed FIO<sub>2</sub>, as applicable; AND
  - 2. oxygen saturations less than or equal to 88% for at least 5 continuous minutes during sleep oximetry (minimum recording time of 2 hours) while the patient is receiving oxygen at 2 LPM or prescribed FIO<sub>2</sub>, whichever is higher; AND
  - 3. OSA and treatment with CPAP has been considered and ruled out.
- C. Continuation of coverage after the 90 day trial period when the following has been demonstrated:
  - 1. Improved AHI and symptom resolution during trial period); and
  - 2. Compliance has been demonstrated as defined by the use of the device for 70% of the nights for an average of 4 or more hours per 24 hour period during a consecutive 30 day period.

VI. Replacement of a positive airway pressure device with an upgraded model will be reviewed for medical necessity and eligible for coverage if:

- A. the patient is compliant with use of the device (*please refer to continuation of coverage after 90 day trial*); and
- B. the device is malfunctioning; and
- C. the device has exceeded the warranty time period; and
- D. required repairs would exceed the cost of a replacement device or the parts that need to be replaced; or
- E. the patient has experienced a change in his or her physiological condition and meets criteria for continued use of the device; or
- F. there has been irreparable change in the device's condition or in a part of the device, due to normal wear and tear.

VII. The monitoring feature/device, stand-alone or integrated, any type, includes all accessories, components and electronics, not otherwise classified (HCPCS: A9279) is considered inclusive to the positive airway pressure device.

VIII. Devices used to clean or sanitize the CPAP or BPAP devices are considered a convenience item and are ineligible for coverage (e.g., SoClean<sup>®</sup>, SoClean 2 Go<sup>®</sup>, SoClean, Inc).

IX. The initiation and management of continuous positive airway pressure ventilation (CPAP) (CPT:94660) is considered **medically appropriate** when performed face-to-face AND criteria for a CPAP device have been met and the device has been approved.

X. Based upon our criteria and assessment of peer-reviewed literature, non-invasive positive pressure ventilators (NIPPV)(e.g., Trilogy100 (Phillips Respironics) (HCPCS: E0466) are considered **medically necessary** for home mechanical ventilation nocturnally (during sleep) for an initial 90 day trial period when

- A. CPAP and BPAP has failed; AND
- B. In adults diagnosed with:

**Medical Policy: POSITIVE AIRWAY PRESSURE DEVICES: CPAP, BiPAP, APAP AND NONINVASIVE POSITIVE PRESSURE VENTILATORS**

**Policy Number: 1.01.06**

**Page: 4 of 10**

1. Neuromuscular diseases; or
2. Thoracic restrictive diseases; or
3. Chronic respiratory failure consequent to chronic obstructive pulmonary disease with:
  - a. arterial blood gas PaCO<sub>2</sub> is greater than or equal to 52 mm Hg while the patient is awake and is breathing either breathing room air or prescribed FIO<sub>2</sub>, as applicable; AND
  - b. oxygen saturations less than or equal to 88% for at least 5 continuous minutes during sleep oximetry (minimum recording time of 2 hours) while the patient is receiving oxygen at 2 LPM or prescribed FIO<sub>2</sub>, whichever is higher; AND
  - c. OSA and treatment with BPAP (without or with back up rate) has been considered and ruled out.
- C. Continuation of coverage after the 90 day trial period when compliance has been demonstrated as defined by the use of the device for 70% of the nights for an average of 4 or more hours per 24 hour period during a consecutive 30 day period during the 90 day trial.
- D. Continuation after one year will be based on compliance that has been demonstrated as defined by the use of the device for 70% of the nights for an average of 4 or more hours per 24 hour period during the most *current* consecutive 30 day period.

*Refer to Corporate Medical Policy # 7.01.41 regarding Surgical Management of Sleep Disorders.*

**POLICY GUIDELINES**

- I. BPAP, CPAP with expiratory relief (e.g., C-Flex technology (Respironics, Inc., Murrysville, PA), or APAP are options for patients who cannot tolerate the high constant air pressure associated with CPAP but wish to continue treatment for OSA.
- II. BPAP devices may or may not include a back up rate. BPAP with a back up rate are timed devices that supply a breath at a specific rate per minute. The device will deliver a breath when the minimum number of breaths per minute has not been met by the user.
- III. Supplemental oxygen and/or humidification can be added to positive pressure devices to increase oxygen saturation and decrease vasomotor rhinitis.
- IV. Prior authorization is contract dependent. Please contact your local Customer (Provider/Member) Services Department to determine contract coverage. Eligibility for reimbursement is based upon the durable medical equipment benefits set forth in the member's subscriber contract.

**DESCRIPTION**

Obstructive sleep apnea (OSA) in an adult is defined as the cessation of airflow through the nose and mouth for at least 10 seconds with a respiratory effort noted. Treatment for OSA is indicated when there is documented sleep related apnea by means of polysomnography or home sleep testing and evidence of clinical impairment such as increased sleepiness or altered cardiopulmonary function.

Obstructive sleep apnea in children is defined as a disorder of breathing during sleep characterized by prolonged partial upper airway obstruction and/or intermittent complete obstruction that disrupts normal ventilation during sleep and normal sleep patterns. In pediatric OSA, the cessation of airflow through the nose and mouth lasts for at least 2 respiratory cycles rather than a time duration of 10 seconds as in an adult. Though adenotonsillectomy is the first-line therapy for pediatric OSA, surgery alone may not be sufficient or may be contraindicated. Continuous positive airway pressure may be an option for these patients. Obstructive hypoventilation (OH) in children is a sleep related breathing disorder that is considered a variation of obstructive sleep apnea. Children with OH may have an AHI within normal ranges, but have episodic periods of hypercarbia as evidenced by elevated measurements in the end-tidal CO<sub>2</sub>.

Central sleep apnea is defined as the cessation of airflow through the nose and mouth for at least 10 seconds with no respiratory effort noted. The cessation in breathing can be caused by problems involving brain mechanisms, or an

## **Medical Policy: POSITIVE AIRWAY PRESSURE DEVICES: CPAP, BiPAP, APAP AND NONINVASIVE POSITIVE PRESSURE VENTILATORS**

**Policy Number: 1.01.06**

**Page: 5 of 10**

obstructive component. Approximately only 4% of patients undergoing PSG in a sleep laboratory are diagnosed with central sleep apnea, making this an uncommon condition.

Mixed sleep apnea is a combination of obstructive and central sleep apnea. Not only does the patient have an obstruction in the airway, but the patient may have a neurological dysfunction or cardiopulmonary as well that contributes to the central apnea component.

*Continuous positive airway pressure, or CPAP*, supplies constant pressure throughout the respiratory cycle by raising the intraluminal upper airway pressure above the positive critical transmural pressure of the pharynx or hypopharynx. The pressure is delivered by a flow generator through either nasal mask or modified nasal prongs in order to keep the upper airway patent resulting in adequate ventilation and arterial oxygenation. The pressure used is determined individually, with a range of 3 to 20 cm. water.

*Bilevel Positive Airway Pressure, or BPAP*, is an airway support system, which provides two different levels of pressure delivered via a mask. There is a higher pressure during inspiration and a lower pressure level during expiration. It is an option for patients who cannot tolerate the high constant air pressure associated with CPAP. Bilevel PAP devices with volume and/or pressure with a backup rate feature are types of ventilators that can be used for many applications (as described below) but have not shown effectiveness for treating OSA.

Bilevel positive pressure airflow is also used in noninvasive ventilation for patients with chronic respiratory failure. Outpatient noninvasive positive pressure ventilation has been used in the following situations:

- I. at night for the management of chronic respiratory failure,
- II. for the long term management of neuromuscular disorders with respiratory involvement,
- III. for patients with respiratory insufficiency due to severe kyphoscoliosis, or
- IV. for the improvement of nighttime desaturation and hypoventilation in patients with chest-wall diseases.

*Auto-or self-titrating positive airway pressure, or APAP*, systems utilize an algorithm that uses a pressure transducer and microprocessor to monitor the airway for vibration patterns and then makes air pressure adjustments based on the incidence of apnea or absence of vibration. APAP devices are also referred to as demand positive airway pressure devices (DPAP).

*CPAP* with expiratory relief (e.g., C-Flex technology (Respironics, Inc., Murrysville, PA) is designed to provide pressure relief during expiration, while maintaining optimal pneumatic splinting for effective therapy. *CPAP* with expiratory relief technology monitors the patient's airflow during expiration and reduces expiratory pressure proportional to expiratory flow. *CPAP* with C-Flex technology can increase compliance in those patients who find it difficult or uncomfortable to breathe out against the continuous positive pressure. This technology can be applied to *CPAP*, *BPAP* and *APAP* devices.

Most PAP machines now contain data cards which have the ability to record and transmit daily use rates of the device. In addition, most PAP machines can also be equipped with a modem (either wired or wireless) that is capable of transmitting data on a daily basis to a manufacturer-owned database. The ability to detect disturbances in a variety of measures, including air flow, amount of time used, and mask leak, is also included in most commercially available PAP machines.

### **RATIONALE**

*CPAP* is effective in the treatment of adult patients with documented obstructive sleep apnea. The symptoms associated with obstructive sleep apnea (OSA) such as excessive daytime sleepiness, impaired cognition, and mood disorders are reduced or eliminated with the consistent use of *CPAP*. *CPAP* is not indicated in individuals with simple snoring that is not associated with pauses in respirations. Treatment recommendations for obstructive sleep apnea are based primarily on the respiratory disturbance index (RDI) or the apnea hypopnea index (AHI), the severity of the presenting symptoms and the existence and severity of co-morbid conditions. (The RDI is defined as the total number of obstructive apneas,

**Medical Policy: POSITIVE AIRWAY PRESSURE DEVICES: CPAP, BiPAP, APAP AND NONINVASIVE POSITIVE PRESSURE VENTILATORS**

**Policy Number: 1.01.06**

**Page: 6 of 10**

hypopneas and respiratory event related arousals per hour; the AHI is defined as the total number of apneas and hypopneas per hour).

Current studies conclude that the extent of improvement in excessive daytime sleepiness is similar between CPAP and APAP. There is no clinical evidence that supports the use of APAP without the results of polysomnography or its use while waiting for polysomnography to be completed.

Comparative studies of CPAP and CPAP-C-Flex have demonstrated similar outcomes in the improvement of OSA. The addition of C-Flex can result in increased comfort and improved compliance in patients who are intolerant of the constant expiratory pressure of traditional CPAP.

Obstructive sleep apnea in children is a common condition that can result in severe complications if left untreated. Complications may include growth abnormalities, neurologic disorders, and cor pulmonale, especially in severe cases. Polysomnography is the best diagnostic technique shown to quantitate the sleep abnormalities associated with sleep disordered breathing, but there is an absence of widely accepted normative data for AHI/RDI in children. Adenotonsillectomy may be a surgical option for children with OSA. For those patients with specific surgical complications, minimal adeno-tonsillar tissue or persistent OSA after adenotonsillectomy, CPAP, observation, medications and additional upper airway surgeries are treatment options.

Positive Airway Pressure has not been approved by the FDA for use in children less than 7 years of age.

**CODES**

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- *CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.*
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*

**CPT Codes**

<b>Code</b>	<b>Description</b>
94660	Continuous positive airway pressure ventilation (CPAP), initiation and management

*Copyright © 2019 American Medical Association, Chicago, IL*

**HCPCS Codes**

<b>Code</b>	<b>Description</b>
A7027-A7039	Accessories/supplies, code range for positive pressure airway devices (code range)
A7044- A7046	Accessories/supplies, code range for positive pressure airway devices (code range)
A9279	Monitoring feature/device, stand-alone or integrated, any type, includes all accessories, components and electronics, not otherwise classified
E0466	Home ventilator, any type, used with noninvasive interface, (e.g., mask, chest shell)
E0470	Respiratory assist device, bi-level pressure capability, without back-up rate feature, used with noninvasive interface, e.g., nasal or facial mask (intermittent assist device with continuous positive airway pressure device)
E0471	Respiratory assist device, bi-level pressure capability, with backup rate feature, used with noninvasive interface, e.g., nasal or facial mask (intermittent assist device with continuous positive airway pressure device)
E0472	Respiratory assist device, bi-level pressure capability, with backup rate feature, used with invasive interface, e.g., tracheostomy tube (intermittent assist device with continuous positive airway pressure device)
E0561	Humidifier, nonheated, used with positive airway pressure device
E0562	Humidifier, heated, used with positive airway pressure device
E0601	Continuous airway pressure (CPAP) device

*Proprietary Information of Univera Healthcare*

**Medical Policy: POSITIVE AIRWAY PRESSURE DEVICES: CPAP, BiPAP, APAP AND NONINVASIVE POSITIVE PRESSURE VENTILATORS**

**Policy Number: 1.01.06**

**Page: 7 of 10**

**ICD10 Codes**

<b>Code</b>	<b>Description</b>
G47.00	Insomnia, unspecified
G47.10	Hypersomnia, unspecified
G47.20	Circadian rhythm sleep disorder, unspecified type
G47.30	Sleep apnea, unspecified
G47.8	Other sleep disorders
G47.9	Sleep disorder, unspecified
J96.00-J96.02	Acute respiratory failure (code range)
J96.90-J96.92	Respiratory failure, unspecified (code range)

**REFERENCES**

- \*Alajmi M, et al. Impact of continuous positive airway pressure therapy on blood pressure in patients with obstructive sleep apnea hypopnea: a meta-analysis of randomized controlled trials. Lung 2007 Mar-Apr;185(2):67-72.
- \*American Academy of Pediatrics. Clinical practice guideline: Diagnosis and management of childhood obstructive sleep apnea syndrome. Pediatrics 2002;109(4):704-712.
- \*American Academy of Sleep Medicine. Practice parameters for the use of auto-titrating continuous positive airway pressure devices for titrating pressures and treating adult patients with obstructive sleep apnea. Sleep 2002;25(2):143-147.
- \*American Thoracic Society. Cardiorespiratory sleep studies in children. Establishment of normative data and polysomnographic predictors of morbidity. Am J Respir Crit Care Med 1999 Oct;160(4):1381-1387.
- Aurora RN, et al. The treatment of central sleep apnea syndromes in adults: practice parameters with an evidence-based literature review and meta-analysis. Sleep 2012 Jan 1;35(1):17-40.
- \*Barnes M, et al. A randomized controlled trial of continuous positive airway pressure in mild obstructive sleep apnea. Am J Respir Crit Care Med 2002 Mar 15;165(6):773-745.
- Bazzano LA, et al. Effect of nocturnal nasal continuous positive airway pressure on blood pressure in obstructive sleep apnea. Hypertension 2007 Aug;50(2):417-423.
- BlueCross BlueShield Association. Diagnosis and Management of Obstructive Sleep Apnea Syndrome. Medical Policy Reference Manual. Policy #2.01.18. 2017 Jun 08.
- Buchner NJ, et al. Continuous positive airway pressure treatment of mild to moderate obstructive sleep apnea reduces cardiovascular risk. Am J Respir Crit Care Med 2007 Dec 15;176(12):1274-1280.
- \*Chervin RD, et al. Inattention, hyperactivity, and symptoms of sleep-disordered breathing. Pediatrics 2002 Mar;109(3):449-456.
- \*Chai CL, et al. Continuous positive airway pressure delivery interfaces for obstructive sleep apnea. Cochrane Database Syst Rev. 2006 Oct 18;(4):CD005308.
- \*Chervin RD, et al. Hyperactivity and polysomnographic findings in children evaluated for sleep-disordered breathing. Sleep 2001 May 1;24(3):313-320.
- DiFeo N, et al. Predictors of positive airway pressure therapy adherence in children: a prospective study. J Clin Sleep Med 2012 Jun 15;8(3):279-286.
- \*Downey R, et al. Nasal continuous positive airway pressure use in children with obstructive sleep apnea younger than 2 years of age. Chest 2000 Jun;117(6):1608-1612.

**Medical Policy: POSITIVE AIRWAY PRESSURE DEVICES: CPAP, BiPAP, APAP AND NONINVASIVE POSITIVE PRESSURE VENTILATORS**

**Policy Number: 1.01.06**

**Page: 8 of 10**

Drager LF, et al. Effects of continuous positive airway pressure on early signs of atherosclerosis in obstructive sleep apnea. Am J Respir Crit Care Med 2007 Oct 1;176(7):706-712.

\*Faccenda JF, et al. Randomized placebo-controlled trial of continuous positive airway pressure on blood pressure in the sleep apnea-hypopnea syndrome. Am J Crit Care Med 2001;163:344-348.

\*Giles TL, et al. Continuous positive airway pressure for obstructive sleep apnea. Cochrane Database Syst Rev 2006 Jul 19;3:CD001106.

Halbower AC, et al. Childhood obstructive sleep-disordered breathing. A clinical update and discussion of technical innovation and challenges. Chest 2007 Dec;132(6):2030-2041.

\*Henke KG, et al. Effect of nasal continuous positive airway pressure on neuropsychological function in sleep apnea-hypopnea syndrome. Am J Crit Care Med 2001;163:911-917.

\*Hudgel DW, et al. A long-term randomized, cross-over comparison of auto-titrating and standard nasal continuous airway pressure. Sleep 2000;23(5):645-648.

\*Kushida CA, et al. Practice parameters for the use of continuous and bilevel positive airway pressure devices to treat adult patients with sleep-related breathing disorders. Sleep 2006 Mar 1;29(3):375-381.

Lam B, et al. Randomized study of three non-surgical treatments in mild to moderate obstructive sleep apnoea. Thorax 2007 Apr;62(4):354-9.

\*Marcus CL, et al. Use of nasal continuous positive airway pressure as treatment of childhood obstructive sleep apnea. J Pediatr 1995 Jul;127(1):88-94.

\*Massa E, et al. The use of nasal continuous positive airway pressure to treat obstructive sleep apnea. Arch Dis Child 2002 Nov;87(5):438-443.

\*McNicholas. Compliance with nasal CPAP therapy for obstructive sleep apnoea: how much is enough? Eur Respir J 1997;10:969-970.

\*Morgenthaler TI, et al. Practice parameters for the medical therapy of obstructive sleep apnea. Sleep 2006 Aug 1;29(8):1031-1035.

Morgenthaler T, et al. Practice parameters for the use of autotitrating continuous positive airway pressure devices for titrating pressures and treating adult patients with obstructive sleep apnea syndrome: an update for 2007. An American Academy of Sleep Medicine report. Sleep 2008 Jan 1;31(1):141-147.

Mulgrew AT, et al. Diagnosis and initial management of obstructive sleep apnea without polysomnography: a randomized validation study. Ann Intern Med 2007 Feb 6;146(3):157-166.

Mulgrew AT, et al. Efficacy and patient satisfaction with autotitrating CPAP with variable expiratory pressure vs standard CPCP: a two-night randomized crossover trial. Sleep Breath 2007 Mar;11(1):31-37.

Murphy PB, et al. Effect of home noninvasive ventilation with oxygen therapy vs oxygen therapy alone on hospital readmission or death after an acute COPD exacerbation: a randomized clinical trial. JAMA 2017;317(21):2177-2186.

\*Nieminen P, et al. Snoring and obstructive sleep apnea in children. A 6-month follow-up study. Arch Otolaryngol Head Neck Surg 2000;126(4):481-486.

Nolan GM, et al. Auto-titrating versus fixed positive pressure therapy in mild to moderate obstructive sleep apnoea. Sleep 2007 Feb 1;30(2):189-194.

Patrino V, et al. Fixed and autoadjusting continuous positive airway pressure treatments are not similar in reducing cardiovascular risk factors in patients with obstructive sleep apnea. Chest 2007 May;131(5):1393-1399.

Peñuelas O, et al. Noninvasive positive-pressure ventilation in acute respiratory failure. CMAJ 2007;177(10):1211-1218.



## **Medical Policy: POSITIVE AIRWAY PRESSURE DEVICES: CPAP, BiPAP, APAP AND NONINVASIVE POSITIVE PRESSURE VENTILATORS**

**Policy Number: 1.01.06**

**Page: 9 of 10**

\*Pieters TH, et al. Acceptance and long-term compliance with nCPAP in patients with obstructive sleep apnoea syndrome. Eur Respir J 1996;9:939-944.

\*Randerath W, et al. Autoadjusting CPAP therapy based on impedance efficacy, compliance and acceptance. Am J Respir Care Med 2001;163:652-657.

\*Sin DD, et al. Can continuous positive airway pressure therapy improve the general health status of patients with obstructive sleep apnea? A clinical effectiveness study. Chest 2002 Nov;122(5):1679-1685.

\*Sin DD, et al. Long-term compliance rates to continuous positive airway pressure in obstructive sleep apnea. A population-based study. Chest 2002 Feb;121(2):430-5.

\*Waters KA, et al. Obstructive sleep apnea: the use of nasal CPAP in 80 children. Am J Respir Crit Care Med 1995 Aug;152(2):780-785.

Weaver TE, et al. Continuous positive airway pressure treatment of sleepy patients with milder obstructive sleep apnea: results of the CPAP Apnea Trial North American Program (CATNAP) randomized clinical trial. Am J Respir Crit Care Med 2012 Oct 1;186(7):677-683.

\*White J, et al. Continuous positive airway pressure for obstructive sleep apnea. Cochrane Database Syst Rev 2002;(2):CD001106.

Windisch W. Noninvasive positive pressure ventilation in COPD. Breathe 2011;8(2):115-123.

\*Veasey SC, et al. Medical therapy for obstructive sleep apnea: a review by the Medical Therapy for Obstructive Sleep Apnea Task Force of the Standards of Practice Committee of the American Academy of Sleep Medicine. Sleep 2006 Aug 1;(29(8):1036-1044.

Zhou L, et al. Home noninvasive positive pressure ventilation with built-in software in stable hypercapnic COPD: a short-term prospective, multicenter, randomized, controlled trial. Int J Chron Obstruct Pulmon Dis 2017 Apr 27;12:1279-1286.

\*Key Article

### **KEY WORDS**

C-Flex, Demand, DPAP, Obstructive sleep apnea, OSA, noninvasive positive pressure ventilation.

### **CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS**

There is currently a Local Coverage Determination (LCD) for Positive Airway Pressure Devices for the Treatment of Obstructive Sleep Apnea. Please refer to the following LCD website for Medicare members:

[https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33718&ver=16&CntrctrSelected=389\\*1&Cntrctr=389&s=41&DocType=Active&bc=AggAAAQBAAA&](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33718&ver=16&CntrctrSelected=389*1&Cntrctr=389&s=41&DocType=Active&bc=AggAAAQBAAA&)

There is currently a Local Coverage Determination (LCD) for Respiratory Assist Devices. Please refer to the following LCD website for Medicare members:

<https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33800&ver=13>

There is currently a National Coverage Determination (NCD) for Durable Medical Equipment. Please refer to the following NCD website for Medicare members: <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?&NCDId=190&ncdver=1&NCDsect=280.1&bc=BEAAAAAAAAQAAAA==&>

**Medical Policy: POSITIVE AIRWAY PRESSURE DEVICES: CPAP, BiPAP, APAP AND NONINVASIVE POSITIVE PRESSURE VENTILATORS**

**Policy Number: 1.01.06**

**Page: 10 of 10**