

# MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	ALLERGY TESTING
Policy Number	2.01.10
Category	Technology Assessment
Effective Date	10/18/01
Revised Date	10/16/02, 10/15/03, 09/16/04, 11/17/05, 09/21/06, 12/20/07, 09/18/08, 09/17/09, 09/16/10, 09/15/11, 09/20/12, 09/19/13, 09/18/14, 09/17/15, 9/15/16, 11/16/17, 01/17/19
Product Disclaimer	<ul style="list-style-type: none"> <li>If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</li> <li>If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit.</li> <li>If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</li> </ul>

## POLICY STATEMENT

- I. Based upon our criteria and review of the peer-reviewed literature, the following tests are **medically appropriate** in the diagnosis of the allergic patient:

CODE	DESCRIPTION	GUIDELINE
95004	Percutaneous tests (scratch, puncture, prick) with allergenic extracts, immediate type reaction, including test interpretation and report, specify number of tests	The number of tests required may vary widely from patient to patient, depending upon the patient's history, and may require up to 70 tests.
95017	Allergy testing, any combination of percutaneous (scratch, puncture, prick) and intracutaneous (intradermal), sequential and incremental, with venoms, immediate type reaction, including test interpretation and report, specify number of tests	Usually used when percutaneous testing is not considered to be sensitive enough to the cause of an allergic reaction. The number of tests required may vary widely from patient to patient, depending upon the patient's history, and may require up to 40 tests.
95018	Allergy testing, any combination of percutaneous (scratch, puncture, prick) and intracutaneous (intradermal), sequential and incremental, with drugs or biologicals, immediate type reaction, including test interpretation and report, specify number of tests	
95024	Intracutaneous (intradermal) tests with allergenic extracts, immediate type reaction, including test interpretation and report, specify number of tests	
95027	Intracutaneous (intradermal) tests, sequential and incremental, with allergenic extracts for airborne allergens, immediate type reaction, including test interpretation and report, specify number of tests	A physician or other qualified health care provider uses intracutaneous tests, sequential and incremental, with allergenic extracts for airborne allergens, immediate type reaction, to determine a patient's specific allergies. The number of tests must be specified ( <i>each sequential test = 1 unit</i> ). This code includes test interpretation and provider report. (serial endpoint testing)

**Medical Policy: ALLERGY TESTING****Policy Number: 2.01.10****Page: 2 of 9**

CODE	DESCRIPTION	GUIDELINE
95028	Intracutaneous (intradermal) tests with allergenic extracts, delayed type reaction, including reading, specify number of tests	Used as a part of an evaluation of the status of immune function. The number of tests is usually small, under 10 tests.
95044	Patch or application test(s) (specify number of tests)	Also known as delayed hypersensitivity testing, this testing modality identifies allergens causing contact dermatitis. The suspected allergens are applied to the patient's back under dressings and allowed to remain in contact with the skin for 48 to 72 hours. The area is then examined for evidence of delayed hypersensitivity reactions.
95052	Photo patch test(s) (specify number of tests)	This test reflects contact photosensitization. A patch of skin is applied with the suspected sensitizer for 48 hours. If no reaction occurs, the area is exposed to a dose of ultraviolet light sufficient to produce inflammatory redness of the skin. If the test is positive, a more severe reaction develops at the patch site than on the surrounding skin.
95056	Photo tests	Photo, or photosensitivity, tests are performed for the evaluation of photosensitivity disorders by irradiating the skin with a specified range of ultraviolet light.
95070	Inhalation bronchial challenge testing (not including necessary pulmonary function tests); with histamine, methacholine, or similar compounds	Histamine or methacholine is used to perform this test when it is necessary to determine if the patient has hyper-responsive airways. Volatile chemicals are used to perform the test when the allergy is encountered in an occupational setting. If dust, ragweed or other common allergens are the suspected cause of the problem, this test is <b>not medically appropriate</b> since skin tests can be used in these situations.
95071	Inhalation bronchial challenge testing (not including necessary pulmonary function tests); with antigens or gases, specify	
95076	Ingestion challenge test (sequential and incremental ingestion of test items, eg, food, drug or other substance); initial 120 minutes of testing	With these tests the patient ingests a food, drug or other substance to which sensitivity is suspected. This may be done in an open or blinded manner. Testing may be done at home, but in some instances of extreme suspected hypersensitivity, it may be performed in the office setting.
95079	Ingestion challenge test (sequential and incremental ingestion of test items, eg, food, drug or other substance); each additional 60 minutes of testing	
82785	Gammaglobulin (immunoglobulin), IgE	Total serum IgE concentration testing is not indicated in most allergic patients, but may be indicated for patients suspected of having allergic bronchopulmonary aspergillosis, immune deficiency disease characterized by increased IgE levels (e.g., Wiskott-Aldrich syndrome, hyper-IgE staphylococcal abscess syndrome), IgE myeloma, pemphigoid, or a poorly controlled moderate to severe asthmatic patient being considered for possible anti IgE treatment.

**Medical Policy: ALLERGY TESTING****Policy Number: 2.01.10****Page: 3 of 9**

CODE	DESCRIPTION	GUIDELINE
86003	Allergen specific IgE; quantitative or semiquantitative, crude allergen extract, each	Commonly known as RAST (radioallergosorbent) testing, these tests detect antigen-specific IgE antibodies in the patient's serum. They are <b>medically appropriate only</b> when testing for allergens (e.g., inhalant, food, insect, drug):
86005	Allergen specific IgE; qualitative, multi-allergen screen (dipstick, paddle or disk)	When direct skin testing is impossible due to extensive dermatitis or marked dermatographism; For patients unable to discontinue use of interfering medications (e.g., antidepressants, antihistamines, or beta blocking agents); For those who have had a near fatal reaction to an allergen;
86008	Allergen specific IgE; quantitative or semiquantitative, recombinant or purified component, each	In children less than four years of age; In patients who will not or cannot cooperate with percutaneous testing due to mental or physical disease (e.g., Down syndrome, mental retardation, dementia);  To follow patients with food allergies and/or insect sting allergies previously documented by history and in-vivo or in-vitro testing;  For patients with suspected latex allergy;  For patients with suspected insect sting allergy in the face of negative skin testing; or  For patients with suspected penicillin allergy.

II. Based upon our criteria and review of the peer-reviewed literature, the following allergy tests have not been medically proven to be effective and are considered **investigational**:

CODE	DESCRIPTION
86001 (E/I)	Allergen specific IgG; quantitative or semiquantitative, each allergen
86343 (E/I)	Leukocyte histamine release test (LHRT)
95060 (E/I)	Ophthalmic mucous membrane test
95065 (E/I)	Direct nasal mucous membrane test
No specific code(s)	Cytotoxicity, Provocative testing (e.g., Rinkel test), Rebeck skin window test

*Refer to Corporate Medical Policy #2.01.04 regarding Clinical Ecology/Multiple Chemical Sensitivities/Idiopathic Environmental Intolerance.*

*Refer to Corporate Medical Policy # 2.01.11 regarding Allergen Immunotherapy.*

*Refer to Corporate Medical Policy # 11.01.03 regarding Experimental and Investigational Services.*

## **Medical Policy: ALLERGY TESTING**

**Policy Number: 2.01.10**

**Page: 4 of 9**

### **POLICY GUIDELINES**

The Federal Employees Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and thus these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

### **DESCRIPTION**

Allergic or hypersensitivity disorders may be manifested by generalized systemic reactions as well as localized reactions in any organ system of the body. The reactions may be acute, subacute or chronic, immediate or delayed and may be caused by numerous offending agents (e.g., pollen, molds, dust, mites, animal dander, stinging insect venoms, foods and drugs).

The optimum management of the allergic patient should include a careful history and physical examination and may include confirming the cause of allergic reaction by information from various testing methods. Once the offending allergenic agent(s) are identified treatment is provided by avoidance, medication and/or immunotherapy.

### **RATIONALE**

Although in vivo (e.g., percutaneous, intracutaneous) testing is presently the preferred method of diagnostic allergy testing for IgE mediated sensitivity in vitro (e.g., RAST) tests are useful when used as stated in the situations identified in the above table.

According to a November 2006 American Academy of Allergy, Asthma and Immunology work group report addressing Allergy Diagnosis in Clinical Practice IgE antibody assay technology has improved with new high binding capacity solid phase matrices, non-isotopic labels for detection antibodies and standards calibrated to the World Health Organization IgE reference preparation. These enhancements have led to an evolution in assay methods from the first generation qualitative assays (e.g., RAST, MAST, EAST), through the second generation semi-quantitative IgE assays (e.g., AutoCAP, Alastat, HYTech, Matrix, MagicLite), to the present state-of-the-art quantitative third generation autoanalyzers. Two third generation immunoassays are the ImmunoCAP System (Phadia) and the Immulite 2000 (Diagnostic Products Corp) whose chemistry is similar to the original RAST, but employ non-isotopic labels and have more rapid throughput with improved precision, accuracy and analytical sensitivity. Their automated chemistries report out allergen-specific IgE antibody quantitatively.

*Serial endpoint testing (SET), or intradermal dilutional testing (IDT), is a form of intradermal skin testing that uses increasing doses of antigen to determine the concentration at which the reaction changes from negative to positive (the "endpoint"). The test has been used for diagnosing allergic disorders and to guide the initiation of immunotherapy by using the endpoint dilution as the starting antigen dose.*

Ferastraoar et al reported (2017) in an independent analysis of 75 patients with over 1600 tests between January 2014 and May 2015 for comparison of skin-prick (SPT), intradermal (IDST), and serum specific immunoglobulin E (ssIgE) testing that: IDST detected more additional environmental sensitizations compared with ssIgE testing. IDST, therefore, may be useful when the SPT and/or ssIgE testing results were negative, but the exposure history indicated relevant allergic sensitization. Serology added only a little more information if both SPT and IDST results were negative but may be useful in combination with SPT if IDST cannot be performed.

In a prospective comparative clinical study (Peltier 2007) 134 subjects were tested for a comparison of intradermal dilutional testing, skin prick testing, and modified quantitative testing for common allergens. The researchers found poor correlation between endpoint and wheal size as graded by a 1 to 4 system and concluded that although a correlation existed, the use of SPT to determine endpoint was inaccurate and dangerous. Modified quantitative testing appears to be a safe alternative to IDT for determining starting doses for immunotherapy. The data supports the safety and efficacy of MQT (combination SPT & IDT)

In a retrospective review of clinical data (random accrual) the authors (Seshul 2006) concluded that IDT is an important step in the determination of the strongest starting dose of immunotherapy that may be safely administered. Initiating immunotherapy in this manner may potentially create significant health care savings by shortening the time required for a patient to reach their individual maximally tolerated dose. The use of a relatively large screening panel is cost effective

## Medical Policy: ALLERGY TESTING

Policy Number: 2.01.10

Page: 5 of 9

and does not increase the average number of antigens treated by immunotherapy. Blended allergy testing techniques that include IDT in their protocol are comparable in cost with commonly used allergy testing protocols. Otolaryngologists often favor IDT (SET) because of its well-documented sensitivity, specificity, safety, and reproducibility. IDT has been compared with many testing modalities used by other physicians to validate the technique as a part of mainstream allergy care.

CDC <https://www.cdc.gov/mmwr/preview/mmwrhtml/rr5511a1.htm> (MMWR 2006) recommends the use of serial endpoint testing (IDT) For patients at high risk for anaphylaxis, including those who 1) have a history of penicillin-related anaphylaxis, asthma, or other diseases that would make anaphylaxis more dangerous or 2) are being treated with beta-adrenergic blocking agents should be tested with 100-fold dilutions of the full-strength skin-test reagents before being tested with full-strength reagents. In these situations, patients should be tested in a monitored setting in which treatment for an anaphylactic reaction is available. If possible, the patient should not have taken antihistamines recently (e.g., chlorpheniramine maleate or terfenadine during the preceding 24 hours, diphenhydramine HCl or hydroxyzine during the preceding 4 days, or astemizole during the preceding 3 weeks).

In a Joint Task Force on Practice Parameters for Drug Allergy; American Academy of Allergy, Asthma and Immunology; American College of Allergy, Asthma and Immunology; Joint Council of Allergy, Asthma and Immunology. (2010) included in their executive summary a statement validating the use of intracutaneous (intradermal) tests are generally used for specific allergens (ie, Hymenoptera venoms and penicillin), but they may also be applied if prick/puncture test results are negative and there is a strong historical likelihood of clinical allergy to specific allergens.

*Leukocyte histamine release testing (LHRT)* is a technique to evaluate the in vitro release of histamine from leukocytes in response to an allergen and provide an in vitro correlate to an in vivo allergic response. Published literature regarding the commercially available LHRTs suffers from not having been performed in a blinded manner or not indicating whether or not there were blinded interpretations of the tests. Some studies included patients with known allergies, which do not represent the same population with equivocal allergy histories that would undergo testing. Studies of LHRT are potentially prone to spectrum, referral, and ascertainment bias, and are not sufficient to permit conclusions on the diagnostic accuracy of the tests. It has been suggested that LHRT may be a valuable test in those patients with discordant results of skin prick testing and RAST testing, but studies focusing on this subgroup of patients have not been identified.

A number of procedures have been shown to be invalid for any clinical purpose. Studies of *cytotoxic tests* and *provocation-neutralization tests* have demonstrated that results are not reproducible. *Electrodermal diagnosis* and *applied kinesiology* have not been evaluated for efficacy. The “*reaginic*” *pulse test* and *chemical analysis of body tissues* have not been substantiated as valid allergy tests. These tests are considered to be investigational.

According to the 2008 American Academy of Allergy, Asthma, and Immunology (AAAAI) and the American College of Allergy, Asthma, and Immunology (ACAAI) joint practice parameter addressing allergy diagnostic testing, IgG and IgG subclass antibody tests for food allergy do not have clinical relevance, are not validated, lack sufficient quality control, and should not be performed. In addition, although a number of investigators have reported modest increases of IgG4 during venom immunotherapy, confirmation and validation of the predictive value of IgG4 for therapeutic efficacy of venom immunotherapy are not yet proven. There is insufficient evidence in the published, peer reviewed scientific literature to support the use of specific IgG antibody testing by RAST or ELISA in the diagnosis or treatment of allergic disease and therefore is investigational.

### CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member’s subscriber contract.*
- *CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.*
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*

#### CPT Codes

Code
------

**Medical Policy: ALLERGY TESTING****Policy Number: 2.01.10****Page: 6 of 9**

<b>Code</b>
Refer to the tables in the policy statement section.

*Copyright © 2019 American Medical Association, Chicago, IL***HCPCS Codes**

<b>Code</b>	<b>Description</b>
No code(s)	

**ICD10 Codes**

<b>Code</b>	<b>Description</b>
B44.0-B44.9	Aspergillosis (code range)
B48.4	Penicillosis
D80.3	Selective deficiency of immunoglobulin G (IgG) subclasses
D82.0	Wiskott-Aldrich syndrome
H10.411- H10.419	Chronic giant papillary conjunctivitis (code range)
H10.45	Other chronic allergic conjunctivitis
J30.0	Vasomotor rhinitis
J30.1-J30.9	Allergic rhinitis (code range)
J45.20-J45.998	Asthma (code range)
L23.0-L23.9	Allergic contact dermatitis (code range)
L24.0-L24.9	Irritant contact dermatitis (code range)
L25.0-L25.9	Unspecified contact dermatitis (code range)
L27.0-L27.9	Dermatitis due to substances taken internally (code range)
L30.0	Nummular dermatitis
L30.2	Cutaneous autosensitization
L30.8	Other specified dermatitis
L30.9	Dermatitis, unspecified
L50.0	Allergic urticaria
L50.3	Dermatographic urticaria
T36.0X5A- T36.0X5S	Adverse effect of penicillins (code range)
T36.1X5A- T36.1X5S	Adverse effect of cephalosporins and other beta-lactam antibiotics (code range)
T39.015A- T39.015S	Adverse effect of aspirin (code range)
T39.095A- T39.095S	Adverse effect of salicylates (code range)
T63.001A- T63.94XS	Toxic effect of contact with venomous animals and plants (code range)
T65.811A- T65.814S	Toxic effect of latex (code range)
T78.00XA- T78.09XS	Anaphylactic reaction due to food (code range)
T78.2xxA	Anaphylactic shock, unspecified, initial encounter

**Medical Policy: ALLERGY TESTING****Policy Number: 2.01.10****Page: 7 of 9**

<b>Code</b>	<b>Description</b>
T78.3xxA	Angioneurotic edema, initial encounter
T78.40XA	Allergy, unspecified, initial encounter
T78.41xA	Arthus phenomenon, initial encounter
T78.49xA	Other allergy, initial encounter
T88.2xxA	Shock due to anesthesia, initial encounter
T88.52XA	Failed moderate sedation during procedure, initial encounter
T88.59xA	Other complications of anesthesia, initial encounter
T88.6XXA	Anaphylactic reaction due to adverse effect of correct drug or medicament properly administered, initial encounter
Z91.010-Z91.09	Allergy status other than drugs & biologicals (code range)

**REFERENCES**

American Academy of Allergy, Asthma and Immunology. Allergy diagnosis in clinical practice. Archived Working Group report. 2006 Dec

[<http://www.aaaai.org/Aaaai/media/MediaLibrary/PDF%20Documents/Practice%20and%20Parameters/Allergy-Diagnosis-2006.pdf>] accessed 10/10/18.

Bernstein DI. Contact Dermatitis for the Practicing Allergist. J Allergy Clin Immunol Pract. 2015 Sep-Oct;3(5):652-8; quiz 659-60. doi: 10.1016/j.jaip.2015.06.006.

\*Bernstein IL, et al. Allergy diagnostic testing: an updated practice parameter. Ann Allergy Asthma Immunol 2008 Mar;100(3 Suppl 3):S1-148.

[<http://www.aaaai.org/Aaaai/media/MediaLibrary/PDF%20Documents/Practice%20and%20Parameters/allergydiagnostictesting.pdf>] accessed 10/8/18.

BlueCross BlueShield Association. Antigen Leukocyte Antibody Test. Medical Policy Reference Manual Policy #2.01.93. 2017 Oct 12.

\*BlueCross BlueShield Association. Leukocyte histamine release test - Archived. Medical Policy Reference Manual Policy #2.04.42. 2009 Dec 12.

BlueCross BlueShield Association. Serial endpoint testing for the diagnosis and treatment of allergic disorders -Archived. Medical Policy Reference Manual Policy #2.01.23. 2013 Jun 13.

\*BlueCross BlueShield Association. Technology Assessment Program (TEC). Serial endpoint testing for the diagnosis and treatment of allergic disorders. 2002 Jul;17(6):1-19.

de Waard-van der Spek FB, et al. EAACI position paper for practical patch testing in allergic contact dermatitis in children. Pediatr Allergy Immunol 2015 Nov;26(7):598-606.

\*Dykewicz MS, et al. Diagnosis and management of rhinitis: complete guidelines of the Joint Task Force on practice parameters in allergy, asthma and immunology. American Academy of Allergy, Asthma and Immunology. Ann Allergy Asthma Immunol 1998 Nov;81(5 Pt 2): 478-518.

Ferastraoar D, et al. Diagnosing environmental allergies: Comparison of skin-prick, intradermal, and serum specific immunoglobulin E testing. Allergy Rhinol (Providence). 2017 Jun 1;8(2):53-62.

Fonacier L. A Practical Guide to Patch Testing. J Allergy Clin Immunol Pract. 2015 Sep-Oct;3(5):669-75.

Fuiano N, et al. Diagnostic performance of the atopy patch test with inhalant allergens. J Investig Allergol Clin Immunol 2015;25(1):34-9.

\*\*Golden DBK, et al. Stinging insect hypersensitivity: A practice parameter update 2016. Ann Allergy Asthma Immunol. 2017 Jan;118(1):28-54.

## **Medical Policy: ALLERGY TESTING**

**Policy Number: 2.01.10**

**Page: 8 of 9**

[<https://www.aaaai.org/Aaaai/media/MediaLibrary/PDF%20Documents/Practice%20and%20Parameters/Stinging-insect-hypersensitivity-2016.pdf>] accessed 10/10/18.

\*Golden DBK, et al. Clinical correlation of the venom-specific IgG antibody level during maintenance venom immunotherapy. J Allergy Clin Immunol 1992 Sep;90(3 Pt 1):386-93.

Johansen JD, et al. European Society of Contact Dermatitis guideline for diagnostic patch testing - recommendations on best practice. Contact Dermatitis 2015 Oct;73(4):195-221.

\*Joint Task Force on Practice Parameters; American Academy of Allergy, Asthma and Immunology; American College of Allergy, Asthma and Immunology; Joint Council of Allergy, Asthma and Immunology. Drug allergy: an updated practice parameter. Ann Allergy Asthma Immunol. 2010 Oct;105(4):259-273.

Marrs T, et al. The diagnosis and management of antibiotic allergy in children: Systematic review to inform a contemporary approach. Arch Dis Child 2015 Jun;100(6):583-8.

Martorell-Aragonés A, et al; Food Allergy Committee of SEICAP (Spanish Society of Pediatric Allergy, Asthma and Clinical Immunology). Position document: IgE-mediated cow's milk allergy. Allergol Immunopathol (Madr) 2015 Sep-Oct;43(5):507-26.

Mowad CM, et al. Allergic contact dermatitis: Patient management and education. J Am Acad Dermatol. 2016 Jun;74(6):1043-54.

\*Peltier J, et al. Comparison of intradermal dilutional testing, skin prick testing, and modified quantitative testing for common allergens. Otolaryngol Head Neck Surg. 2007 Aug;137(2):246-9.

Peters AT, et al; Joint Task Force on Practice Parameters, representing the American Academy of Allergy, Asthma and Immunology, the American College of Allergy, Asthma and Immunology, and the Joint Council of Allergy, Asthma and Immunology. Diagnosis and management of rhinosinusitis: a practice parameter update. Ann Allergy Asthma Immunol 2014 Oct;113(4):347-85.

Rosenfeld RM, et al. Clinical practice guideline (update): adult sinusitis. Otolaryngol Head Neck Surg 2015 Apr;152(2 Suppl):S1-S39.

Seidman MD, et al; Guideline Otolaryngology Development Group. AAO-HNSF. Clinical practice guideline: Allergic rhinitis. Otolaryngol Head Neck Surg 2015 Feb;152(1 Suppl):S1-43.

\*Seshul M, et al, Use of Intradermal Dilutional Testing and Skin Prick Testing: Clinical Relevance and Cost Efficiency Laryngoscope. 2006 Sep;116(9):1530-8.

\*Key Article

### **KEY WORDS**

Allergy tests: Allergen specific IgE, Allergen specific IgG, Challenge, Cytotoxic, Dipstick, Disk, Intracutaneous, Intradermal, Leukocyte histamine release, Mucous membrane, Paddle, Percutaneous, Phadiatop, Prick, Provocation-neutralization, RAST, Rinkel, Scratch, Serial endpoint titration, Skin test.

### **CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS**

There is currently a National Coverage Determination (NCD) addressing Food Allergy Testing and Treatment and Cytotoxic Food Tests. There is also a Local Coverage Determination (LCD) addressing RAST Type Tests. Please refer to the following websites for Medicare Members:

NCD:

[https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=266&ncdver=1&CoverageSelection=Both&ArticleType=All&PolicyType=Final&s=New+York+-+Upstate&Keyword=allergy+testing&KeywordLookUp=Title&KeywordSearchType=And&ncd\\_id=110.11&ncd\\_version=1&basket=ncd%25253A110%25252E11%25253A1%25253AFood+Allergy+Testing+and+Treatment&bc=gAAAABAAAA&](https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=266&ncdver=1&CoverageSelection=Both&ArticleType=All&PolicyType=Final&s=New+York+-+Upstate&Keyword=allergy+testing&KeywordLookUp=Title&KeywordSearchType=And&ncd_id=110.11&ncd_version=1&basket=ncd%25253A110%25252E11%25253A1%25253AFood+Allergy+Testing+and+Treatment&bc=gAAAABAAAA&)



**Medical Policy: ALLERGY TESTING**

**Policy Number: 2.01.10**

**Page: 9 of 9**

[https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=161&ncdver=1&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCA%7cCAL%7cNCD%7cMEDCAC%7cTA%7cMCD&ArticleType=SAD%7cEd&PolicyType=Both&s=41&Keyword=allergy&KeyWordLookUp=Doc&KeyWordSearchType=Exact&kq=true&bc=IAAAACAAAA&](https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=161&ncdver=1&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCA%7cCAL%7cNCD%7cMEDCAC%7cTA%7cMCD&ArticleType=SAD%7cEd&PolicyType=Both&s=41&Keyword=allergy&KeyWordLookUp=Doc&KeyWordSearchType=Exact&kq=true&bc=IAAAACAAAA&bc=IAAAACAAAA&)

**LCD:**

[https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33591&ver=15&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCA%7cCAL%7cNCD%7cMEDCAC%7cTA%7cMCD&ArticleType=SAD%7cEd&PolicyType=Both&s=41&Keyword=allergy&KeyWordLookUp=Doc&KeyWordSearchType=Exact&kq=true&bc=IAAAACAAAA&](https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=33591&ver=15&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCA%7cCAL%7cNCD%7cMEDCAC%7cTA%7cMCD&ArticleType=SAD%7cEd&PolicyType=Both&s=41&Keyword=allergy&KeyWordLookUp=Doc&KeyWordSearchType=Exact&kq=true&bc=IAAAACAAAA&bc=IAAAACAAAA&)