

# MEDICAL POLICY



MEDICAL POLICY DETAILS	
Medical Policy Title	CT (Computed Tomography) Perfusion Imaging of the Brain
Policy Number	6.01.37
Category	Technology Assessment
Original Effective Date	12/18/08
Committee Approval Date	12/17/09, 03/17/11, 05/19/11, 05/24/12, 05/23/13, 05/22/14, 06/18/15, 02/18/16, 02/16/17, 02/15/18, 02/21/19, 02/20/20, 01/21/21, 01/20/22, 01/19/23
Current Effective Date	01/19/23
Archived Date	N/A
Archive Review Date	N/A
Product Disclaimer	<ul style="list-style-type: none"> <li>• If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply.</li> <li>• If a commercial product (including an Essential Plan or Child Health Plus product), medical policy criteria apply to the benefit.</li> <li>• If a Medicaid product covers a specific service, and there are no New York State Medicaid guidelines (eMedNY) criteria, medical policy criteria apply to the benefit.</li> <li>• If a Medicare product (including Medicare HMO-Dual Special Needs Program (DSNP) product) covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.</li> <li>• If a Medicare HMO-Dual Special Needs Program (DSNP) product DOES NOT cover a specific service, please refer to the Medicaid Product coverage line.</li> </ul>

## POLICY STATEMENT

- I. Based upon our criteria and assessment of the peer-reviewed literature, computed tomography (CT) perfusion imaging for assessing patients suspected of having an acute stroke less than twenty-four hours earlier or in triaging stroke patients for interventional thrombectomy or thrombolysis treatment is considered **medically appropriate**.
- II. Based upon our criteria and assessment of the peer-reviewed literature, CT perfusion imaging has not been medically proven to be effective and, therefore, is considered **investigational** for all other indications.

*Refer to Corporate Medical Policy #11.01.03 Experimental or Investigational Services.*

## DESCRIPTION

Perfusion imaging using CT is proposed to provide detailed study of cerebral blood flow (CBF), which may assist in identification of ischemic regions of the brain, especially within the first few hours of an acute stroke. The technique requires either a diffusible inert gas indicator such as xenon (Xe) or a non-diffusible indicator such as an iodinated contrast agent. The CT scanner is then used to capture images as the agent accumulates in cerebral tissues. The technique tracks transient attenuation changes in the blood vessels and brain parenchyma during the first pass of an intravenously injected contrast medium. Results of CT perfusion studies allow calculation of regional cerebral blood volume (CBV), mean transit time (MTT), and CBF. Proposed advantages of CT perfusion imaging are that it is less invasive than angiography and more widely available than magnetic resonance imaging (MRI).

Three CT perfusion imaging approaches use different data acquisition and analysis methods. Whole brain CT perfused blood volume is assessed by a helical scan through the whole brain with and without contrast. First pass perfusion CT (bolus tracking CT perfusion study) acquires repeated images at the same location through a volume of interest during bolus injection and passage of contrast through the region of interest. Dynamic perfusion CT acquires a temporal set of

## **Medical Policy: CT (COMPUTED TOMOGRAPHY) PERFUSION IMAGING OF THE BRAIN**

**Policy Number: 6.01.37**

**Page: 2 of 5**

images through an extended volume of interest (imaging of tissue beyond the absolute width of the detector array) during a bolus injection of contrast.

### **RATIONALE**

Several post-processing software packages have received Section 510(k) marketing clearance from the U.S. Food and Drug Administration (FDA) for use with a CT system, to perform perfusion imaging, e.g., the GE Medical Systems CT Perfusion 4 (March 2006), the Philips Medical Systems Brain Perfusion Option (Feb. 2004) and the Siemens Medical Solutions syngo Perfusion-CT (Dec. 2003).

The American College of Radiology (ACR) and American Society of Neuroradiology (ASNR) published an ACR/ASNR practice guideline (October 2012) for performance of CT perfusion in neuroradiologic imaging. The guidelines state that brain primary indications for perfusion CT in neuroradiology include, but are not limited to: differentiating salvageable ischemic penumbra from unsalvageable ischemic core, distinguishing benign oligemia from true “at-risk” ischemic penumbra, helping identify patients most likely to benefit from thrombolysis or thrombectomy, predicting hemorrhagic transformation in acute ischemic stroke, and identifying patients with malignant profiles, suspected vasospasm-related cerebral ischemia and infarction and/or delayed cerebral ischemia (DCI) following aneurysmal subarachnoid hemorrhage, and cerebral hemorrhage with secondary local ischemia. Brain secondary indications are: (1) follow-up of acute cerebral ischemia or infarction and/or reperfusion in the subacute or chronic phase of recovery, to assist in planning and evaluating the effectiveness of therapy for cervical or intracranial arterial occlusive disease (as an isolated test or in combination with a cerebrovascular reserve challenge) and/or chronic cerebral ischemia; (2) identifying cerebral hyperperfusion syndrome following revascularization; (3) detection of crossed cerebellar diaschisis in acute ischemic stroke; and (3) contrast delay as a predictor of new incident infarct. CT perfusion scanning may also be helpful in the setting of acute traumatic brain injury, the setting of acute seizures, the assessment of neoplastic disease, and in patients with contraindication to MRI-based perfusion imaging or with devices or material in or close to the field of view that would result in nondiagnostic MRI scans. Head and neck primary indications include evaluation of the vascular status of solid tumors where MRI is degraded due to susceptibility artifact from air-containing spaces or from surgical clips or dental work. Head and neck secondary indications include follow-up of tumor response to therapy.

The American Heart Association (AHA)/American Stroke Association (ASA)/American Academy of Neurology (AAN) published guidelines for Early Management of Adults with Ischemic Stroke in January 2013. The “Early Diagnosis, Brain and Vascular Imaging” section of the guideline states that the AHA, ASA, and AAN agree that emergency, non-contrast-enhanced CT scanning of the brain accurately identifies most cases of intracranial hemorrhage and helps distinguish nonvascular causes of neurological symptoms, such as a brain tumor. That section also states that recent technological advances have led to increased interest in more sophisticated, multimodal approaches to acute stroke imaging. Multimodal CT approaches may include non-contrast CT, perfusion CT, and CT angiography (CTA) studies. Whole brain perfusion CT provides a map of cerebral blood volume (CBV) and regions of hypoattenuation on these CBV maps represent the ischemic core. Although this technique has the advantage of providing whole-brain coverage, it is limited by its inability to provide measures of cerebral blood flow (CBF) or mean transit time (MTT). Dynamic perfusion CT has the potential to provide absolute measures of CBF, MTT, and CBV. This technique is limited to four to eight brain slices and provides incomplete visualization of all pertinent vascular territories, although newer (320-slice) CT machines can provide whole-brain coverage with this technique.

The AHA also concluded in the published guidelines for Early Management of Adults with Ischemic Stroke in January 2013 that the usefulness of vascular imaging for predicting responses to treatment before intravenous administration of thrombolytic agents has not been demonstrated. AHA Class I recommendations state that multimodal CT and MR may provide additional information that will improve diagnosis of ischemic stroke. Class II recommendations state that CT perfusion and MRI perfusion and diffusion imaging, including measures of infarct core and penumbra, may be considered for the selection of patients for acute reperfusion therapy beyond the time windows for intravenous fibrinolysis. These techniques provide additional information that may improve diagnosis, mechanism, and severity of ischemic stroke and allow more informed clinical decision-making.

In 2019, the AHA and ASA revised their 2018 guideline statement on the use of CTP for the Early Management of Adults with Ischemic Stroke as follows:

## Medical Policy: CT (COMPUTED TOMOGRAPHY) PERFUSION IMAGING OF THE BRAIN

Policy Number: 6.01.37

Page: 3 of 5

RECOMMENDATION	SOR	LOB	LOE
In patients eligible for IV alteplase, because benefit of therapy is time dependent, treatment should be initiated as quickly as possible and not delayed for additional multimodal neuroimaging, such as CT and MRI perfusion imaging.	I	Strong benefit	B-NR (nonrandomized)
When selecting patients with acute ischemic stroke within 6 to 24 hours of last known normal who have large vessel occlusion in the anterior circulation, obtaining CTP or DW-MRI, with or without MRI perfusion, is recommended to aid in patient selection for mechanical thrombectomy, but only when patients meet other eligibility criteria from one of the RCTs that showed benefit from mechanical thrombectomy in this extended time window.	I	Strong benefit	A (high-quality evidence from multiple RCTs)
In selected patients with acute ischemic stroke (>16 to 24 hours of last normal) and large vessel occlusion, DAWN criteria (which may include imaging findings from CTP) may be used for clinical decision making regarding mechanical thrombectomy.	Ila	Moderate benefit	B-R (nonrandomized)

*CT: computed tomography; CTP: computed tomography perfusion; DW-MRI: diffusion-weighted magnetic resonance imaging; IV: intravenous; LOB: level of benefit; LOE: level of evidence; MRI: magnetic resonance imaging; RCT: randomized controlled trial; SOR: strength of recommendation.*

### Perfusion CT in Brain Tumors

The standard for tumor grading is a histopathologic assessment of tissue. Limitations of histologic assessment include sampling error due to regional heterogeneity and interobserver variation. These limitations can result in inaccurate classification and grading of gliomas. Because malignant brain tumors are characterized by neovascularity and increased angiogenic activity, perfusion imaging has been proposed as a method to assess tumor grade and prognosis. Dynamic contrast-enhanced MRI (DCE-MRI) is the preferred technique, because there is no radiation exposure and a good signal-to-noise ratio. Perfusion CT may be an alternative choice for glioblastoma patients with DCE-MRI examination contraindications. Potential advantages, compared with MR perfusion, include the wider availability, faster scanning times, and lower cost. CTP imaging may also be used to distinguish recurrent tumor from radiation necrosis.

## CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- ***CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.***
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*
- *Code Key: Experimental/Investigational = (E/I), Not medically necessary/ appropriate = (NMN).*

### **CPT Codes**

Code	Description
0042T	Cerebral perfusion analysis using computed tomography with contrast administration, including post-processing of parametric maps with determination of cerebral flood flow, cerebral blood volume, and mean transit time.

*Copyright © 2023 American Medical Association, Chicago, IL*

### **HCPCS Codes**

Code	Description
No code	

## Medical Policy: CT (COMPUTED TOMOGRAPHY) PERFUSION IMAGING OF THE BRAIN

Policy Number: 6.01.37

Page: 4 of 5

### ICD10 Codes

Code	Description
I63.00-I63.9	Cerebral infarction (code range)

### REFERENCES

\*Adams HP Jr, et al. Guidelines for the early management of adults with ischemic stroke: a guideline from the American Heart Association/American Stroke Association Stroke Council, Clinical Cardiology Council, Cardiovascular Radiology and Intervention Council, and the Atherosclerotic Peripheral Vascular Disease and Quality of Care Outcomes in Research Interdisciplinary Working Groups. Circulation 2007 May 22;115(20):e478-534.

Adebayo OD and Culpán G. Diagnostic accuracy of computed tomography perfusion in the prediction of haemorrhagic transformation and patient outcome in acute ischaemic stroke: A systematic review and meta-analysis. European Stroke Journal 2020;5(1):4-16.

\*Agency for Healthcare Research and Quality. Acute stroke: evaluation and treatment. Evidence report-technology assessment (summary) 2005;(127):1-7. [<https://archive.ahrq.gov/clinic/epcs/sums/acstrokesum.htm>] accessed 11/18/22.

\*Allmendinger AM, et al. Imaging of stroke: Part 1, Perfusion CT--overview of imaging technique, interpretation pearls, and common pitfalls. AJR Am J Roentgenol 2012 Jan;198(1):52-62.

\*American College of Radiology (ACR) and the American Society of Neuroradiology (ASNR). Practice guideline for the performance of CT perfusion in neuroradiologic imaging. Resolution 18. 2017 Oct [<https://www.acr.org/-/media/ACR/Files/Practice-Parameters/ct-perfusion.pdf>] accessed 11/18/22.

Austein F, et al. Value of perfusion CT in the prediction of intracerebral hemorrhage after endovascular treatment. Stroke Res Treat 2021 Jul 22;2021:9933015.

Becks MJ, et al. Brain CT perfusion improves intracranial vessel occlusion detection on CT angiography. Journal of Neuroradiology 2019;46:124-129.

\*Chalouhi N, et al. CT perfusion-guided versus time-guided mechanical recanalization in acute ischemic stroke patients. Clin Neurol Neurosurg 2013 Dec;115(12):2471-5.

Chu Y, et al. Diagnostic accuracy of using Alberta Stroke Program Early Computed Tomography Score on CT perfusion map to predict a target mismatch in patients with acute ischemic stroke. Neuroradiology 2022;64:1321-1330.

\*Connolly ES, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. Stroke 2012 Jun;43(6):1711-37.

\*Donahue J, et al. Perfusion CT and acute stroke imaging: foundations, applications, and literature review. J Neuroradiol 2015 Feb;42(1):21-9.

\*González RG, et al. The Massachusetts General Hospital acute stroke imaging algorithm: an experience and evidence based approach. J Neurointerv Surg 2013 May;5(Suppl 1):i7-12.

\*Greenberg ED, et al. Diagnostic accuracy of CT angiography and CT perfusion for cerebral vasospasm: a meta-analysis. AJNR AM J Neuroradiol 2010 Nov;31(10):1853-60.

\*Kelly A, et al. Predictors of rapid brain imaging in acute stroke: analysis of the Get with The Guidelines-Stroke Program. Stroke 2012 May;43(5):1279-84.

\*Kilpatrick MM, et al. CT-based assessment of acute stroke: CT, CT angiography, and xenon-enhanced CT cerebral blood flow. Stroke 2001 Nov;32(11):2543-9.

Klug J, et al. Integrating regional perfusion CT information to improve prediction of infarction after stroke. J Cereb Blood Flow Metab 2021 Mar;41(3): 502-510.

## **Medical Policy: CT (COMPUTED TOMOGRAPHY) PERFUSION IMAGING OF THE BRAIN**

**Policy Number: 6.01.37**

**Page: 5 of 5**

Kuribara T, et al. Ischemic tolerance evaluated by computed tomography perfusion during balloon test occlusion. Journal of Stroke and Cerebrovascular Diseases 2020 Jun;29(6):104807.

McDonough R, et al. Computed tomography-based triage of extensive baseline infarction: ASPECTS and collaterals versus perfusion imaging for outcome prediction. J NeuroInterv Surg 2021;13:869-874.

\*Obach V, et al. Multimodal CT-assisted thrombolysis in patients with acute stroke: a cohort study. Stroke 2011 Apr;42(4):1129-31.

Onishi S, et al. Perfusion computed tomography parameters are useful for differentiating glioblastoma, lymphoma, and metastasis. World Neurosurg 2018;119:e890-e897.

\*Prabhakaran S, et al. Perfusion-based selection for endovascular reperfusion therapy in anterior circulation acute ischemic stroke. AJNR Am J Neuroradiol 2014 Jul;35(7):1303-8.

\*Sheth KN, et al. Advanced modality imaging evaluation in acute ischemic stroke may lead to delayed endovascular reperfusion therapy without improvement in clinical outcomes. J Neurointerv Surg 2013; 5 Suppl 1:i62-5.

Tsurukiri J, et al. Thrombectomy for stroke at 6-24 hours without perfusion CT software for patient selection. Journal of Stroke and Cerebrovascular Diseases, 2019 Mar; 28(3):774-781.

\*Turk AS, et al. CT perfusion-guided patient selection for endovascular recanalization in acute ischemic stroke: a multicenter study. J Neurointerv Surg 2013 Nov;5(6):523-7.

\*Weist R, et al. Detection of regional blood perfusion changes in epileptic seizures with dynamic brain perfusion CT - a pilot study. Epilepsy Res 2006 Dec;72(2-3):102-10.

\*Wintermark M, et al. Comparison of admission perfusion computed tomography and qualitative diffusion- and perfusion-weighted magnetic resonance imaging in acute stroke patients. Stroke 2002 Aug;33(8):2025-31.

\*Wintermark M, et al. Prognostic accuracy of cerebral blood flow measurement by perfusion computed tomography, at the time of emergency room admission, in acute stroke patients. Ann Neurol 2002 Apr;51(4):417-32.

\*Key Article

### **KEY WORDS**

Dynamic Perfusion CT, Multimodal CT, PCT, Perfusion CT, Xenon-enhanced CT, XeCT.

### **CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS**

There is currently a Local Coverage Determination (LCD) for Computed Tomography Cerebral Perfusion Analysis. Please refer to the following LCD website for Medicare members:

<https://www.cms.gov/medicare-coverage-database/view/lcd.aspx?LCDId=38667&ver=3>