



MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	Gender Reassignment/Gender Affirming Surgery and Treatments
Policy Number	7.01.84
Category	Contract Clarification
Effective Date	10/28/10
Revised Date	12/08/11, 10/25/12, 10/24/13, 10/23/14, 12/10/15, 12/8/16, 04/26/18, 04/24/19, 04/23/20
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan product) covers a specific service, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

- I. Based on our criteria and assessment of peer-reviewed literature, bilateral mastectomy, with or without chest reconstruction, including nipple/areola reconstruction and tattooing, for individuals who were assigned female at birth and are transitioning to male, has been shown to be a beneficial and effective intervention for gender dysphoria, and, therefore, is considered **medically appropriate** when **ALL** of the following criteria are met:
- A. The patient has received a letter of referral from a qualified mental health professional (*refer to Policy Guidelines below*); and
 - B. The patient has been diagnosed with persistent gender dysphoria, including all of the following:
 1. The desire to live and be accepted as a member of the identified gender, usually accompanied by the wish to make their body as congruent as possible with the preferred gender through surgery and hormone treatment; and
 2. The gender dysphoria has been present persistently for at least one year; and
 3. The condition is not a symptom of another mental disorder or a chromosomal abnormality; and
 4. The condition causes clinically significant distress or impairment in social, occupational, or other important areas of functioning; and
 - C. The patient has the capacity to make a fully informed decision and to consent to treatment, as well as the ability to comply with all aftercare instructions, including recommended medical, surgical, nursing, and/or psychological care recommended by the individual's providers; and
 - D. The patient has reached the age of majority (18 years of age or older), or, if under the age of majority, meets all of the following criteria for early intervention:
 1. has consent from parent(s)/guardian(s) for surgery; and
 2. has identified as transgender for at least two years; and
 3. has been living in the desired gender role for at least one year; and
 4. has been receiving testosterone treatment for at least one year; and
 5. has received an additional letter of referral from a second qualified mental health professional or physician (*refer to Policy Guidelines below*); and
 6. has compelling reasons impacting their physical and/or psychological well-being, as documented by the patient's mental health/adolescent medicine provider(s); and
 7. any significant medical or mental health concerns that are present are reasonably well controlled.

Note: Hormone treatment history is not required for adults seeking chest reconstruction (including mastectomy) surgery.

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS

Policy Number: 7.01.84

Page: 2 of 13

- II. Based on our criteria and assessment of peer-reviewed literature, breast augmentation/implants, including nipple/areola reconstruction and tattooing, for individuals who were assigned male at birth and are transitioning to female has been shown to be a beneficial and effective intervention for gender dysphoria, and, therefore, is considered **medically appropriate** when **ALL** of all the following criteria are met:
- A. The patient has received a recommendation letter from a qualified mental health professional (*refer to Policy Guidelines below*); and
 - B. The patient has been diagnosed with persistent gender dysphoria, including all of the following:
 - 1. The desire to live and be accepted as a member of the opposite sex, usually accompanied by the wish to make his or her body as congruent as possible with the preferred gender through surgery and hormone treatment; and
 - 2. The gender dysphoria has been present persistently for at least one year; and
 - 3. The condition is not a symptom of another mental disorder or a chromosomal abnormality; and
 - 4. The condition causes clinically significant distress or impairment in social, occupational, or other important areas of functioning; and
 - C. The patient has the capacity to make a fully informed decision and to consent to treatment, as well as the ability to comply with all aftercare instructions, including recommended medical, surgical, nursing, and/or psychological care recommended by the individual's providers; and
 - D. The patient has reached the age of majority (18 years of age or older); and
 - E. If significant medical or mental health concerns are present, they must be reasonably well controlled; and
 - F. The patient has completed a minimum of 24 months of hormone therapy, unless hormone therapy is medically contraindicated, or the patient is otherwise unable to take hormones.
- III. Based on our criteria and assessment of peer-reviewed literature, gonadectomy (i.e., hysterectomy and oophorectomy in a birth-assigned female transitioning to male and orchiectomy in a birth-assigned male transitioning to female) has been shown to be effective and, therefore, is considered **medically appropriate** when all the following criteria are met:
- A. The patient has received two recommendation letters submitted by qualified mental health professionals, or has received one letter from a qualified mental health professional and one letter from a physician (MD, DO), as follows:
 - 1. One letter should be submitted by a mental health professional with whom the individual has had ongoing interactions sufficient to:
 - a. establish a diagnosis of severe and persistent gender dysphoria;
 - b. rule out other diagnoses that might confound the diagnosis of gender dysphoria;
 - c. identify pertinent patient strengths, stressors, and supports; and
 - d. diagnose and address other relevant psychological disorders that might otherwise interfere with the individual's success.
 - 2. The second mental health professional or physician providing a recommendation is not required to have an ongoing relationship with the individual, but should have significant experience assessing individuals with gender dysphoria and/or evaluating decision making capacity in individuals prior to major medical procedures and surgeries (*refer to Policy Guidelines below for additional information*); and
 - B. The patient has an established and well-documented history of gender dysphoria, diagnosed by a mental health professional and present for a minimum of one year, including all of the following:
 - 1. Distress with their assigned gender and with the physical attributes or secondary sex characteristics of their assigned gender;
 - 2. A desire to do away with current secondary sexual characteristics and/or a desire to change their secondary sex characteristics in order to bring them more in line with their internally experienced gender;
 - 3. Noticeable gender distress that causes clinically significant impairment in social, occupational, or other areas of functioning; and
 - 4. Distress and associated symptoms are not better explained by another psychological disorder or by a chromosomal abnormality or intersex condition; and
 - C. The patient has the capacity to make a fully informed decision and to consent to treatment; and

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS

Policy Number: 7.01.84

Page: 3 of 13

- D. The patient has reached the age of majority (18 years or older); and
 - E. If significant medical or mental health conditions are present, the patient has appropriate medical and psychiatric providers in place, symptoms are under reasonably good control, and a plan for continued follow-up of these conditions is in place; and
 - F. The patient has a history of 12 months of continuous hormone therapy consistent with the patient's gender goals, unless hormone therapy is medically contraindicated or the patient has a history of a severe medical or psychiatric adverse effect from hormonal treatments).
- IV. Based on our criteria and assessment of peer-reviewed literature, genital reconstructive surgery (i.e., vaginectomy, urethroplasty, metoidioplasty, phalloplasty, scrotoplasty, and placement of a testicular prosthesis and erectile prosthesis in individuals transitioning from female to male; penectomy, vaginoplasty, labiaplasty, and clitoroplasty in individuals transitioning from male to female) has been medically proven to be effective and, therefore, is considered **medically appropriate** when **ALL** of the following criteria are met:
- A. The patient has received two recommendation letters submitted by qualified mental health professionals or has received one letter from a qualified mental health professional and one letter from a physician (MD, DO), as follows:
 - 1. One letter should be submitted by a mental health professional with whom the individual has had ongoing interactions sufficient to:
 - a. establish a diagnosis of severe and persistent gender dysphoria;
 - b. rule out other diagnoses that might confound the diagnosis of gender dysphoria;
 - c. identify pertinent patient strengths, stressors, and supports; and
 - d. diagnose and address other relevant psychological disorders that might otherwise interfere with the individual's success.
 - 2. The second mental health professional or physician providing a recommendation is not required to have an ongoing relationship with the individual, but should have significant experience assessing individuals with gender dysphoria and/or evaluating decision making capacity in individuals prior to major medical procedures and surgeries (*refer to Policy Guidelines below for additional information*).
 - B. The patient has an established and well-documented history of gender dysphoria, diagnosed by a mental health professional, including all the following characteristics:
 - 1. Distress with their assigned gender and with the physical attributes or secondary sex characteristics of their assigned gender;
 - 2. A desire to do away with current secondary sexual characteristics and/or a desire to change their secondary sex characteristics in order to bring them more in line with their internally experienced gender;
 - 3. Noticeable gender distress that causes clinically significant impairment in social, occupational, or other areas of functioning; and
 - 4. Distress and associated symptoms are not better explained by another psychological disorder or by a chromosomal abnormality or intersex condition; and
 - C. Patient has the capacity to make a fully informed decision and to consent to treatment (*refer to Policy Guidelines*); and
 - D. Patient has reached the age of majority (age 18 years and older); and
 - E. If significant medical or mental health conditions are present, the patient has appropriate medical and psychiatric providers in place, symptoms are under reasonably good control, and a plan for continued follow-up of these conditions is in place; and
 - F. The patient has a history of 12 months of continuous hormone therapy consistent with the patient's gender goals, unless hormone therapy is medically contraindicated or the patient has history of a severe medical or psychiatric adverse effect from hormonal treatments).
- V. Surgical/non-surgical interventions that may be elected by some transgender individuals and that are generally considered **not medically necessary** as part of gender reassignment/gender affirming treatment include:
- A. Abdominoplasty;
 - B. Blepharoplasty;
 - C. Liposuction;

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS

Policy Number: 7.01.84

Page: 4 of 13

- D. Rhinoplasty;
 - E. Face lift;
 - F. Facial bone reconstruction/facial feminization surgery;
 - G. Jaw shortening/sculpturing;
 - H. Chin/nose implants;
 - I. Voice modification surgery;
 - J. Voice therapy/lessons;
 - K. Tracheal shaving/thyroid chondroplasty;
 - L. Hair removal, hair transplantation, electrolysis, or hairplasty, unless required for vaginoplasty or phalloplasty;
 - M. Removal of redundant skin;
 - N. Skin resurfacing;
 - O. Prosthetic or filler substances to alter contour;
 - P. Mastopexy;
 - Q. Calf and pectoral implants; and
 - R. Lip reduction/enhancement.
- VI. Services to reverse gender reassignment/gender affirming surgery are considered **not medically necessary** and, therefore, **not eligible for coverage**.
- VII. Based on our criteria and the lack of peer-reviewed literature, histrelin acetate subcutaneous implant is considered **investigational** for suppression of puberty in transgender individuals.

Refer to Corporate Medical Policy #3.01.15 Behavioral Health Treatment for Gender Dysphoria.

Refer to Corporate Medical Policy #4.01.05 Assisted Reproductive Technologies for Infertility.

Refer to Corporate Medical Policy #7.01.55 Blepharoplasty with or without Levator Muscle Advancement.

Refer to Corporate Medical Policy #7.01.11 Cosmetic and Reconstructive Procedures.

Refer to Corporate Medical Policy #7.01.53 Abdominoplasty and Panniculectomy.

Refer to Corporate Medical Policy #8.01.13 Speech Pathology/Therapy for voice therapy requests.

Refer to Corporate Medical Policy #10.01.01 Breast Reconstruction Surgery for nipple/areola reconstruction and tattooing.

Refer to Corporate Medical Policy #11.01.26 Sex Specific Services for Transgender Individuals.

POLICY GUIDELINES

- I. Coverage of oral or self-administered hormone therapy and hormone inhibitor agents is dependent upon the prescription drug benefits of the member's subscriber contract.
- II. Two state-licensed health care professionals must recommend gender reassignment/gender affirming genital surgery. One professional must be a qualified mental health provider with whom the patient has an established and ongoing professional relationship. The recommendation must specify the provider's competency in transgender care. The second professional may be a psychiatrist, psychologist, licensed clinical social worker, or physician. The providers must establish that gender affirming surgery is medically necessary to treat the individual's gender dysphoria and that the individual demonstrates full capacity for informed decision making, consent, and compliance. Capacity includes: an understanding of common risks and complications, short- and long-term outcomes (e.g., effects on sexual function/fertility), options available to address fertility or sexual function concerns, and the expected benefits associated with surgery. Further, informed decision making requires that an individual have realistic expectations from surgical treatment and have the ability to plan for and comply with the recommendations of their providers with regard to surgical, medical, nursing, and psychological care following surgery. Based on a comprehensive assessment of capacity, the mental health provider should attest to the individual's readiness and appropriateness for the surgery being proposed (note: if breast/chest surgery is the only procedure being requested in an adult patient, only one

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS

Policy Number: 7.01.84

Page: 5 of 13

mental health provider recommendation is required; however, this recommendation must come from a mental health provider with whom the individual has an established and ongoing professional relationship, and must include a comprehensive assessment of capacity as outlined above).

- III. An established and ongoing professional relationship is defined as one in which the provider has had ongoing interactions with the individual sufficient to:
 - A. establish a diagnosis of severe and persistent gender dysphoria;
 - B. rule out other diagnoses that might confound the diagnosis of gender dysphoria;
 - C. identify pertinent patient strengths, stressors, and supports; and
 - D. diagnose and address other relevant psychological disorders that might otherwise interfere with the individual's success.
- IV. For individuals with considerable comorbidities or a history of severe symptoms (due to gender dysphoria, minority stress, or other mental conditions), the provider may provide a recommendation for surgery that includes an appropriate treatment plan for addressing and mitigating these symptoms, stressors, or conditions in the pre-and post-surgical periods.
- V. The patient should have sufficient medical, nursing, and emotional support to adequately address needs in the post-operative, recovery, and healing period. (For individuals having surgery remotely but returning home less than two weeks following surgery, the patient must have medical providers in place who will be following the surgery both in the home area and/or in the city where surgery is to be performed.)
- VI. In-home medical/nursing supports are required in the post-operative period, which may include family members, partners, or friends; or, if no family member, partner, or friend is involved, sufficient alternative options for aftercare support (e.g., visiting nurse, etc.) must be identified.
- VII. If the patient is to have surgery in an out-of-town location and return home, the medical and/or surgical providers who will be responsible for the patient's post-surgical care and who will manage any complications should be identified. (Note: A plan to use urgent care/ emergency care is not sufficient.)
- VIII. The Health Plan recognizes that treatments and services to address gender dysphoria remain limited. In some areas, especially those areas remote from larger cities, finding surgical and/or mental health providers may be more challenging. For this reason, many individuals elect to have gender affirming treatments and surgeries in remote locations where more comprehensive services and providers with more experience are available. Individuals are encouraged to utilize these centers and facilities for mental health assessments and supports, in addition to surgical treatments. Assessments done by mental health providers at such facilities will be considered carefully. Further, for individuals experiencing difficulties in locating medical, surgical, or mental health providers for treatment of gender dysphoria and/or for support in pursuing gender affirming treatment, care management services are available free of charge to members through the Health Plan.

DESCRIPTION

Gender dysphoria, previously known as gender identity disorder (GID), involves a conflict between an individual's gender as perceived (or assigned) and the individual's own internal experience of their gender. Gender dysphoria as defined in the Diagnostic and Statistical Manual of Mental Disorders, 5th Edition (DSM-5) requires that there be a noticeable difference between an individual's assigned gender and their gender identity, which is present for at least six months and which causes clinically significant impairments in the individual's functioning. Gender dysphoria is not equivalent to gender non-conformity, gender expansiveness, or to the term "transgender." Not all transgender individuals experience gender dysphoria, although many do. Gender dysphoria occurs when the individual feels significant discomfort and a desire to change their gender socially and/or physically. In addition, the individual may feel an intense need to transform their gender and/or may have severe difficulty coping with their conditions. People with gender dysphoria may report a feeling of being born the wrong sex. The causes of gender dysphoria and the developmental factors associated with it are not well-understood. Gender affirming surgical options in order to assist an individual to transition to a gender consistent with their identity are now well-established and effective interventions for the treatment of extreme cases of gender dysphoria for those with sufficient preparation and readiness. Gender reassignment therapy is an umbrella term for all

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS

Policy Number: 7.01.84

Page: 6 of 13

procedures regarding gender reassignment and usually consists of a real-life experience in the desired role, hormone replacement therapy to modify secondary sex characteristics, and gender reassignment surgery to alter primary sex characteristics. This therapeutic approach is sometimes labeled triadic therapy due to the three key elements involved. Individuals with gender dysphoria require psychological treatment long before reassignment therapy begins and usually continue it permanently after the “transition.”

Gender reassignment or gender affirming surgery is a permanent change to a patient’s sexual identity and is not reversible. Therefore, a careful and accurate diagnosis is essential for treatment and can be made only as part of a long-term diagnostic process involving a multidisciplinary specialty approach that includes an extensive case history; gynecological, endocrinological and urological examination; and a clinical psychiatric/psychological examination. The goal of gender reassignment or gender affirming surgery is to align the individual’s physical appearance and genital anatomy with his/her gender identity. Gender reassignment or gender affirming surgery involves a series of procedures that will make male genitals into female genitals or vice versa (e.g., penectomy, orchiectomy, vaginoplasty, hysterectomy, salpingo-oophorectomy, colpectomy, metoidioplasty, phalloplasty) and will reshape a male body into a body with female appearances or vice versa (e.g., mastectomy, facial feminization surgery, nose/chin implants, jaw sculpturing, tracheal shaving, voice modification surgery, hair removal).

Gender dysphoria is a DSM-5-recognized medical condition and a pre-requisite for gender-affirming surgery coverage. Many individuals seek mental health treatment to address gender dysphoria, however gender-affirming treatment (including surgery) is recognized as effective in treating gender dysphoria. At the same time, gender transition is a stressful experience for most individuals, and this is especially true in the post-operative period.

RATIONALE

A diagnosis of gender dysphoria is based on the DSM criteria. The DSM-5 provides for one overarching diagnosis of gender dysphoria, with separate specific criteria for children and for adolescents and adults. In adolescents and adults, gender dysphoria diagnosis involves a difference between one’s experienced/expressed gender and assigned gender, and significant distress or problems functioning. It lasts at least six months and is shown by at least two of the following:

- I. A marked incongruence between one’s experienced/expressed gender and primary and/or secondary sex characteristics;
- II. A strong desire to be rid of one’s primary and/or secondary sex characteristics;
- III. A strong desire for the primary and/or secondary sex characteristics of the other gender;
- IV. A strong desire to be of the other gender;
- V. A strong desire to be treated as the other gender; or
- VI. A strong conviction that one has the typical feelings and reactions of the other gender.

Psychological techniques that attempt to treat gender dysphoria via attempts to alter the individual’s gender identity or expression to one considered appropriate for the person’s assigned sex (conversion treatments) have typically been shown to be ineffective. Most providers agree (and research supports) that the most effective and reasonable course of treatment for individuals with gender dysphoria is gender transition, which, for many, will eventually involve gender affirming surgery of some type; and, for individuals with persistent gender dysphoria, this option is considered medically necessary. This conclusion is supported by evidence that individuals with untreated gender dysphoria have higher rates of depression, anxiety, substance abuse problems, and suicide.

The literature related to gender reassignment surgery has numerous limitations (e.g., lack of controlled studies, evidence not collected prospectively, large number of patients lost to follow-up). However, the majority of patients in case series and cohort studies experienced successful outcomes in terms of subjective self-assessment of their surgery, as well as low rates of regret.

The World Professional Association for Transgender Health or WPATH (formerly known as the Harry Benjamin International Gender Dysphoria Association) Standards of Care (SOC) for the Health of Transsexual, Transgender, and Gender Nonconforming People and the DSM-5 criteria are widely accepted as definitive documents in the area of gender dysphoria treatment. Per WPATH, the rationale for a preoperative, 12-month experience of living in an identity-congruent gender role is as follows: The criterion noted for some types of genital surgeries—i.e., that patients engage in

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS

Policy Number: 7.01.84

Page: 7 of 13

12 continuous months of living in a gender role that is congruent with their gender identity—is based on expert clinical consensus that this experience provides ample opportunity for patients to experience and socially adjust in their desired gender role, before undergoing irreversible surgery. The social aspects of changing one’s gender role are usually challenging—often more so than the physical aspects. Changing gender role can have profound personal and social consequences, and the decision to do so should include an awareness of what the familial, interpersonal, educational, vocational, economic, and legal challenges are likely to be, so that people can function successfully in their gender role. Support from a qualified mental health professional and from peers can be invaluable in ensuring a successful gender role adaptation (Bockting, 2008). The duration of 12 months allows for a range of different life experiences and events that may occur throughout the year (e.g., family events, holidays, vacations, season-specific work or school experiences). During this time, patients should present consistently, on a day-to-day basis and across all settings of life, in their desired gender role. This includes coming out to partners, family, friends, and community members (e.g., at school, work, other settings).

The criteria in the SOC are supported by evidence-based, peer-reviewed journal publications. Several studies have shown that extensive long-term trials of hormonal therapy and real-life experience living as the other gender, as well as social support and acceptance by peer and family groups, greatly improve psychological outcomes in patients undergoing gender reassignment surgery (Eldh, 1997; Landen, 1998). A study reported by Monstrey and colleagues (2001) described the importance of close cooperation between the many medical and behavioral specialties required for proper treatment of patients with gender dysphoria who wish to undergo gender reassignment surgery.

One study of 188 patients undergoing gender reassignment surgery found that dissatisfaction with surgery was highly associated with sexual preference, psychological co-morbidity, and poor pre-operative body image and satisfaction (Smith, 2005). Other researchers (MI Lobato, *et al.* (2006), JC Goodard, *et al.* (2007)) reported good overall cosmetic results and high patient satisfaction in studies related to the early and long-term follow-up of patients undergoing gender reassignment surgery (n=19 and n=233, respectively).

Gender affirming surgeries present significant medical and psychological risks, and involve long-term, often-irreversible results. Further, gender transition is a highly stressful process for most, in many cases because of the stress placed upon transgender individuals by others in their families, communities, and work sites, and within society. Many individuals who experience gender dysphoria do benefit from psychological support, if only to allow them a safe environment in which to explore their own minority-stress experience, and to process and plan for a transition that is individualized, safe, and affirming for them. In most cases, a step-wise approach to gender affirming transition interventions is prudent. In adults for whom secondary sex characteristics are established, a careful approach to transition and to gender affirming treatment allows for accurate diagnosis and long-term treatment planning by a multidisciplinary team that includes behavioral, medical and surgical specialists. Both short-term and long-term outcomes are improved in individuals’ whose transitions have proceeded according to careful plan and for whom multidisciplinary services and supports have been put in place. As with the treatment of any condition for which mental health symptoms are present, a thorough psychological analysis by a qualified practitioner is of fundamental importance. Once a diagnosis of gender dysphoria has been established, a trial of hormone therapy is an evidenced-based and helpful treatment intervention which is generally prescribed prior to embarking upon more invasive surgical treatment options. In addition, careful consideration of realistic, safe, and acceptable “real-life” or social transition experiences may be of help for many individuals who are planning for or receiving gender affirming treatment.

Histrelin acetate is a gonadotropin releasing hormone (GnRH) agonist that is FDA-approved for the treatment of children with central precocious puberty or for palliative treatment of advanced prostate cancer. There is insufficient evidence in the literature to support the use of histrelin in the treatment of gender dysphoria or for the suppression of puberty onset in transgender youth.

CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member’s subscriber contract.*
- ***CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY STATEMENTS AND GUIDELINES CAREFULLY.***

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS**Policy Number: 7.01.84****Page: 8 of 13**

- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.

CPT Codes

Code	Description
11920	Tattooing, intradermal introduction of insoluble opaque pigments to correct color defects of skin, including micropigmentation; 6.0 sq cm or less
11921	6.1 to 20.0 sq cm
11922	each additional 20.0 sq cm, or part thereof (List separately in addition to code for primary procedure)
19303	Mastectomy, simple, complete
19318	Reduction mammoplasty
19324	Mammoplasty, augmentation; without prosthetic implant
19325	with prosthetic implant
19350	Nipple/areola reconstruction
53410	Urethroplasty, 1-stage reconstruction of male anterior urethra
53415	Urethroplasty, transpubic or perineal, 1-stage, for reconstruction or repair of prostatic or membranous urethra
53420	Urethroplasty, 2-stage reconstruction or repair of prostatic or membranous urethra; first stage
53425	second stage
53430	Urethroplasty, reconstruction of female urethra
54125	Amputation penis, complete
54400	Insertion of penile prosthesis; non-inflatable (semi-rigid)
54401	Insertion of penile prosthesis; inflatable (self-contained)
54405	Insertion of multi-component, inflatable penile prosthesis, including placement of pump, cylinders, and reservoir
54520	Orchiectomy, simple (including subcapsular), with or without testicular prosthesis, scrotal or inguinal approach
54660	Insertion testicular prosthesis (separate procedure)
54690	Laparoscopy, surgical, orchiectomy
55175	Scrotoplasty; simple
55180	Scrotoplasty, complicated
55970	Intersex surgery, male to female
55980	Intersex surgery, female to male
56625	Vulvectomy simple; complete
56800	Plastic repair introitus
56805	Clitoroplasty for intersex state
57110	Vaginectomy, complete removal of vaginal wall;
57111	Vaginectomy, complete removal of vaginal wall; with removal of paravaginal tissue (radical vaginectomy)
57291	Construction of artificial vagina; without graft
57292	with graft
57335	Vaginoplasty for intersex state

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS

Policy Number: 7.01.84

Page: 9 of 13

Code	Description
58150	Total abdominal hysterectomy, (corpus and cervix), with or without removal of tube(s), with or without removal of ovary(s)
58180	Supracervical abdominal hysterectomy (subtotal hysterectomy), with or without removal of tube(s), with or without removal of ovary(s)
58260	Vaginal hysterectomy, for uterus 250 g or less;
58262	for uterus 250 g or less; with removal of tube(s), and/or ovary(s)
58275	with total or partial vaginectomy;
58290	for uterus greater than 250 g;
58291	for uterus greater than 250 g; with removal of tube(s) and/or ovary(s)
58541	Laparoscopy, surgical, supracervical hysterectomy, for uterus 250 g or less;
58542	for uterus 250 g or less; with removal of tube(s) and/or ovary(s)
58543	for uterus greater than 250 g;
58544	for uterus greater than 250 g; with removal of tube(s) and/or ovary(s)
58550-58554	Laparoscopy, surgical, with vaginal hysterectomy, with or without removal of ovaries and/or tubes (code range)
58570-58573	Laparoscopy, surgical, total hysterectomy with or without removal of ovaries and/or tubes (code range)
58661	Laparoscopy, surgical; with removal of adnexal structures (partial or total oophorectomy and/or salpingectomy)
58700	Salpingectomy, complete or partial, unilateral or bilateral (separate procedure)
58720	Salpingo-oophorectomy, complete or partial, unilateral or bilateral (separate procedure)
58940	Oophorectomy, partial or total, unilateral or bilateral;
58953	Bilateral salpingo-oophorectomy with omentectomy, total abdominal hysterectomy and radical dissection for debulking;
58999	Unlisted procedure, female genital system (nonobstetrical)

Copyright © 2020 American Medical Association, Chicago, IL

HCPC Codes

Code	Description
J9225 (E/I)	Histrelin implant (Vantas), 50 mg
J9226 (E/I)	Histrelin implant (Supprelin LA), 50 mg

ICD10 Codes

Code	Description
F64.0-F64.9	Gender identity disorders (code range)
Z87.890	Personal history of sex reassignment

REFERENCES

Previously titled Sex Reassignment Surgery.

American Academy of Child and Adolescent Psychiatry. Practice parameters on gay, lesbian, or bisexual orientation, gender nonconformity, and gender discordance in children and adolescents. *J Am Acad Child Adolesc Psychiatry* 2012 Sep;51(9):957-74.

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS

Policy Number: 7.01.84

Page: 10 of 13

American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. Arlington, VA: American Psychiatric Publishing, 2013.

American Psychological Association. Guidelines for psychological practice with transgender and gender nonconforming people. 2015 Dec;70(9):832-864.

Andréasson M, et al. Patient-reported outcome measures used in gender confirmation surgery: a systematic review. Plast Reconstr Surg 2018 Apr;141(4):1026-1039.

Aydin D, et al. Transgender surgery in Denmark from 1994 to 2015: 20-year follow-up study. J Sex Med 2016 April;13(4):720-725.

*Bradley SJ and Zucker KJ. Gender identity disorder: A review of the past 10 years. J Am Acad Child Adolesc Psychiatry 1997;36(7):872-80.

Byne W, et al. Report of the American Psychiatric Association Task Force on treatment of gender identity disorder. Arch Sex Behav 2012 Aug;41(4):759-96.

Castellano E, et al. Quality of life and hormones after sex reassignment surgery. J Endocrinol Invest 2015 Dec;38(12):1373-1381.

*Cohen-Kettenis PT, et al. Transsexualism: a review of etiology, diagnosis and treatment. J Psychosom Res 1999; 46(4):315-333.

Cohen WA, et al. Female-to-male transgender chest contouring: A systematic review of outcomes and knowledge gaps. Ann Plast Surg 2019 Nov;83(5):589-593.

*Committee on Adolescent Health Care. Committee opinion no. 685: care for transgender adolescents. Obstet Gynecol 2017 Jan;129(1):e11-e16.

*Committee on Health Care for Underserved Women. Committee Opinion no.512. Health care for transgender individuals. Obstet Gynecol 2011 Dec;118(6):1454-8.

de Blok CJM, et al. Breast development in transwomen after 1 year of cross-sex hormone therapy: results of a prospective multicenter study. J Clin Endocrinol Metab 2018 Feb 1;103(2):532-538.

Dy GW, et al. Presenting complications to a reconstructive urologist after masculinizing genital reconstructive surgery. Urology 2019 Oct;132:202-206.

*Eldh J, et al. Long-term follow up after sex reassignment surgery. Scand J Plast Reconstr Surg Hand Surg 1997;31(1):39-45.

*Goddard JC, et al. Feminizing genitoplasty in adult transsexuals: early and long-term surgical results. BJU Int 2007 Sep;100(3):607-13.

*Gooren LJ. Clinical practice. Care of transsexual persons. NEJM 2011 Mar 31;364(13):1251-7.

Gorin-Lazard A, et al. Hormonal therapy is associated with better self-esteem, mood, and quality of life in transsexuals. J Nerve Ment Dis 2013 Nov;201(11):996-1000.

*Hembree WC, et al. Endocrine treatment of transsexual persons: an Endocrine Society clinical practice guideline. J Clin Endocrinol Metab 2009 Sep;94(9):3132-54.

*Hembree WC, et al. Endocrine treatment of gender-dysphoric/gender-incongruent persons: an Endocrine Society clinical practice guideline. J Clin Endocrinol Metab 2017 Nov;102(11):1-35.
[<https://academic.oup.com/jcem/article/102/11/3869/4157558>] accessed 3/6/20.

*Hepp U, et al. A. Psychiatric comorbidity in gender identity disorder. J Psychosom Res 2005;58(3):259-61.

Heylens G, et al. Effects of different steps in gender reassignment therapy on psychopathology: a prospective study of persons with a gender identity disorder. J Sex Med 2014 Jan;11(1):19-26.

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS

Policy Number: 7.01.84

Page: 11 of 13

*Klein C, et al. Sexual functioning in transsexuals following hormone therapy and genital surgery: a review. J Sex Med 2009 Nov;6(11):2922-39.

*Kuhn A, et al. Vaginal prolapse. Pelvic floor function, and related symptoms 16 years after sex reassignment surgery in transsexual. Fertil Steril 2011 Jun;95(7):2379-82.

*Landen M, et al. Factors predictive of regret in sex reassignment. Acta Psychiatr Scand 1998;97(4):284-9.

*Lawrence AA. Factors associated with satisfaction or regret following male-to-female sex reassignment surgery. Arch Sex Behav 2003 Aug;32(4):299-315.

*Lawrence AA. Sexuality before and after male-to-female sex reassignment surgery. Arch Sex Behav 2005 Apr;34(2):147-66.

Levy JA, et al. Male-to-female gender reassignment surgery: An institutional analysis of outcomes, short-term complications, and risk factors for 240 patients undergoing penile-inversion vaginoplasty. Urology 2019 Sep;131:228-233.

Lindqvist EK, et al. Quality of life improves early after gender reassignment surgery in transgender women. Eur J Plast Surg 2017;40(3):223-226.

*Lobato MI, et al. Follow-up of sex reassignment surgery in transsexuals: a Brazilian cohort. Arch Sex Behav 2006 Dec;35(6):711-5.

Lopez CM, et al. Trends in the use of puberty blockers among transgender children in the United States. J Pediatr Endocrinol Metab 2018 Jun 27;31(6):665-670.

Manrique OJ, et al. Complications and patient-reported outcomes in male-to-female vaginoplasty-where we are today: a systematic review and meta-analysis. Ann Plast Surg 2018 Jun;80(6):684-691.

Massie JP. Predictors of patient satisfaction and postoperative complications in penile inversion vaginoplasty. Plast Reconstr Surg 2018 Jun;141(6):911e-921e.

*Meriggiola MC, et al. Endocrine treatment of transsexual persons: an Endocrine Society clinical practice guideline: commentary from a European perspective. Eur J Endocrinol 2010 May;162(5):831-3.

*Midence K and Hargreaves I. Psychosocial adjustment in male-to-female transsexuals: an overview of the research evidence. J Psychol 1997;131(6):602-14.

*Monstrey S, et al. Surgical therapy in transsexual patients: a multi-disciplinary approach. Acta Chir Belg 2001;101(5):200-9.

Ngaage LM, et al. Health insurance coverage of gender-affirming top surgery in the United States. Plast Reconstr Surg 2019 Oct;144(4):824-833.

Olson-Kennedy, J, et al. Chest reconstruction and chest dysphoria in transmasculine minors and young adults: comparisons of nonsurgical and postsurgical cohorts. JAMA Pediatr 2018 May 1;172(5):431-436.

Papadopoulos NA, et al. Male-to-female sex reassignment surgery using the combined technique leads to increased quality of life in a prospective study. Plast Reconstr Surg 2017 Aug;140(2):286-294.

Rafferty J, et al. Ensuring comprehensive care and support for transgender and gender-diverse children and adolescents. Pediatrics 2018 Oct;142(4) [<http://pediatrics.aappublications.org/content/142/4/e20182162>] accessed 3/6/20.

*Schlatterer K, et al. Multistep treatment concept of transsexual patients. Exp Clin Endocrin Diab 1996;104(6):413-9.

*Selvaggi G, et al. Gender identity disorder: general overview and surgical treatment for vaginoplasty in male-to-female transsexuals. Plast Reconstr Surg 2005;116(6):135e-45e.

Selvaggi G, et al. The 2011 WPATH standards of care and penile reconstruction in female-to-male transsexual individuals. Adv Urol 2012;58:1712.

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS

Policy Number: 7.01.84

Page: 12 of 13

- *Smith YL, et al. Sex reassignment: outcomes and predictors of treatment for adolescent and adult transsexuals. Psychol Med 2005;35(1):89-99.
- *Smith YL, et al. Adolescents with gender identity disorder who were accepted or rejected for sex reassignment surgery: a prospective follow-up study. J Am Acad Child Adolesc Psychiatry 2001 Apr;40(4):472-81.
- *Sohn M, et al. Gender identity disorders: diagnostic and surgical aspects. J Sex Med 2007 Sep;4(5):1193-207.
- *Sutcliffe PA, et al. Evaluation of surgical procedures for sex reassignment: a systematic review. J Plast Reconstr Aesthet Surg 2009 Mar;62 (3):294-306.
- Van der Sluis WB, et al. Long-term follow-up of transgender women after secondary intestinal vaginoplasty. J Sex Med 2016 April;13(4):702-710.
- Weigert R, et al. Patient satisfaction with breasts and psychosocial, sexual, and physical well-being after breast augmentation in male-to-female transsexuals. Plast Reconstr Surg 2013 Dec;132(6):1421-9.
- Weinforth G, et al. Quality of life following male-to-female sex reassignment surgery. Dtsch Arztebl Int 2019 Apr 12;116(15):253-260.
- Wernick JA, et al. A systematic review of the psychological benefits of gender-affirming surgery. Urol Clin North Am 2019 Nov;46(4):475-486.
- *Wierckx K, et al. Quality of life and sexual health after sex reassignment surgery in transsexual men. J Sex Med 2011 Dec;8(12):3379-88.
- *World Professional Association for Transgender Health (WPATH). Standards of Care for the Health of Transsexual, Transgender, and Gender Nonconforming people. 2012 7th version.
[https://www.wpath.org/media/cms/Documents/SOC%20v7/SOC%20V7_English.pdf] accessed 3/6/20.
- *Key Article

KEY WORDS

Gender dysphoria, Gender identity disorder, GID, gender reassignment surgery, genital correction surgery, genital reassignment surgery, genital reconstruction, gender realignment surgery, gender confirmation surgery, intersex, transsexualism, transsexual surgery.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently a National Coverage Determination (NCD) for Gender Dysphoria and Gender Reassignment Surgery. Please refer to the following NCD website for Medicare Members. <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCID=368&ncdver=1&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCA%7cCAL%7cNCD%7cMEDCAC%7cTA%7cMCD&ArticleType=SAD%7cEd&PolicyType=Both&s=41&Keyword=gender+dysphoria&KeywordLookUp=Doc&KeywordSearchType=Exact&kq=true&bc=IAAAACAAAA&>

A final decision memo was issued in August 2016 by CMS for Gender Dysphoria and Gender Reassignment Surgery. The memo is located at: <https://www.cms.gov/medicare-coverage-database/details/ncd-a-decision-memo.aspx?NCAID=282&CoverageSelection=National&Keyword=gender+reassignment+surgery&KeywordLookUp=Title&KeywordSearchType=And&bc=gAAAACAACAAAA%3d%3d&>

Per CMS Manual, Pub 100-03, Medicare National Coverage Determinations, Transmittal 194, change request 9981 was issued 03/03/17 with implementation 04/04/2017. Effective for claims with dates of service on or after August 30, 2016, coverage determinations for gender reassignment surgery, under section 1862(a)(1)(A) of the Social Security Act and any other relevant statutory requirements, will continue to be made by the local Medicare Administrative. Contractors (MACs)

Medical Policy: GENDER REASSIGNMENT/GENDER AFFIRMING SURGERY AND TREATMENTS

Policy Number: 7.01.84

Page: 13 of 13

on a case-by-case basis. This transmittal is located at: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Transmittals/2017Downloads/R194NCD.pdf>