

MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	HER-2 TESTING IN INVASIVE BREAST OR GASTRIC CANCER USING FLUORESCENCE IN SITU HYBRIDIZATION (FISH) OR IMMUNOHISTOCHEMISTRY (IHC) ASSAYS
Policy Number	2.02.31
Category	Laboratory Test
Effective Date	07/20/06
Revised Date	10/18/07, 10/23/08, 10/29/09, 10/28/10, 03/17/11, 02/16/12, 01/17/13, 01/16/14, 01/22/15, 03/17/16, 05/18/17, 03/15/18, 03/21/19, 3/19/20
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

- I. Based upon our criteria and review of the peer-reviewed literature, testing of breast cancer biopsy samples for determination of HER-2/neu gene status in newly diagnosed or newly metastatic invasive breast cancer has been medically proven to be effective and, therefore, is considered **medically appropriate** when one of the following methods are used:
 - A. Fluorescence in situ hybridization (FISH) testing; or
 - B. Immunohistochemistry (IHC) testing.
- II. Based upon our criteria and review of the peer-reviewed literature, testing for determination of HER-2/neu gene status is considered **medically appropriate** prior to initiation of treatment with the monoclonal antibody Trastuzumab (Herceptin®) in patients with locally advanced, recurrent, or metastatic adenocarcinoma of the stomach or gastroesophageal (GE) junction adenocarcinoma who have not received prior treatment for metastatic disease.
- III. Based upon our criteria and review of the peer-reviewed literature, validation of an equivocal (e.g. inconclusive) HER-2 result via FISH or IHC testing is considered **medically appropriate** as follows:
 - A. IHC score of 2+ should be subjected to reflex testing by a validated complementary (e.g. FISH) method;
 - B. FISH result of an average HER2 gene copy should be a number of greater than four to less than six, or a gene amplification ratio of 1.8 to 2.2, or should undergo:
 1. Counting of additional cells; or
 2. Retesting by FISH; or
 3. Reflex testing by a validated IHC method.

Refer to Corporate Medical Policy #2.02.35 Molecular Analysis for Targeted Therapy of Non-Small Cell Lung Cancer

Refer to Corporate Medical Policy #2.02.41 Genotyping- RAS Mutation Analysis in Metastatic Colorectal Cancer

POLICY GUIDELINES

- I. According to the American Society of Clinical Oncology and College of American Pathologists (ASCO/CAP) guidelines:
 - A. A positive HER-2 result is:
 1. IHC score of 3+; or

Medical Policy: HER-2 TESTING IN INVASIVE BREAST OR GASTRIC CANCER USING FLUORESCENCE IN SITU HYBRIDIZATION (FISH) OR IMMUNOHISTOCHEMISTRY (IHC) ASSAYS

Policy Number: 2.02.31

Page: 2 of 8

2. FISH result of greater than or equal to 6.0 HER-2 gene copies per nucleus or dual-probe HER2/CEP17 ratio greater than or equal to 2.0.
- B. A negative HER-2 result is:
 1. IHC score of 0 or 1+; or
 2. FISH result of less than 4.0 HER-2 gene copies per nucleus or a dual-probe *HER2/CEP17* ratio of less than 2.0.
- C. An equivocal (e.g. inconclusive) HER-2 result is:
 1. IHC score of 2+; or
 3. FISH result with a dual-probe *HER2/CEP17* ratio of less than 2.0 and an average HER2 copy number of greater than 4.0 and fewer than 6.0 gene copies per nucleus.

II. This policy does not address the use of Herceptin® in the treatment of breast cancer.

DESCRIPTION

The human epidermal growth factor receptor 2 (HER-2/neu) oncogene is found in high concentrations on the cell surfaces of various cancers. Each cell should have only two copies of the HER-2/neu gene. If a cell has more than two copies, it produces too much of the HER-2/protein. Excessive amounts of this protein (overexpression) causes cells to reproduce uncontrollably. Although the HER-2/neu protein is overexpressed in several epithelial cancers including breast, ovarian, thyroid, lung, salivary gland, stomach, colon and prostate cancer, it has been reported that approximately one-third of breast cancer patients overexpress HER-2/neu. This overexpression appears to be associated with more aggressive disease and usually results from amplification (multiple extra copies) of the HER-2/neu gene in malignant cells.

HER-2/neu testing is used to assist with the selection of breast cancer candidates for treatment with the monoclonal antibody Trastuzumab (Herceptin®). It has been proven by multiple Phase III studies that patients with HER-2/neu overexpressing breast cancer will benefit from therapy with Herceptin®, which may be indicated either as first line therapy or in combination with chemotherapy. Additionally, knowing a tumor's HER-2/neu level may give information about the nature of the cancer and the expected outcome, which can help in the selection of appropriate treatment.

HER-2/neu testing is used to assist with the selection of gastric cancer candidates for treatment with the monoclonal antibody Trastuzumab (Herceptin®). As reported in the NCCN 2011 Gastric Cancer Practice Guidelines, the ToGA study was the first randomized, prospective, multicenter, phase III trial to evaluate the efficacy and safety of trastuzumab in HER2-positive gastric cancer in combination with cisplatin and a fluoropyrimidine. The results of this study confirmed that trastuzumab plus standard chemotherapy is superior to chemotherapy alone in patients with HER2-positive advanced gastric cancer. In this study, 594 patients with HER2-positive gastroesophageal (GE) and gastric adenocarcinoma (locally advanced, recurrent, or metastatic) were randomized to receive trastuzumab plus chemotherapy (5-fluorouracil or capecitabine and cisplatin) or chemotherapy alone. There was a significant improvement in the median overall survival with the addition of trastuzumab to chemotherapy compared to chemotherapy alone (13.5 vs. 11.1 months, respectively.) Safety profiles were similar with no unexpected adverse events in the trastuzumab. There was also no difference in symptomatic congestive heart failure between the two arms. This establishes trastuzumab plus chemotherapy as a new standard of care for the treatment of patients with a HER2-expressing advanced gastric and GE cancers.

Two distinct methods are used for detection of the HER-2/neu oncogene in breast and gastric cancer patients. Both methods can be performed on archived or current specimens.

Immunohistochemistry (IHC) is performed on breast tumor tissue removed at surgery and measures protein expression of the HER-2/neu gene. Results are reported as a range from 0 to 3+. 1+ is negative, 2+ is indeterminate or weak (may be positive or negative), and 3+ is positive. An example of an IHC HER-2/neu test is the HercepTest®.

The *fluorescence in situ hybridization* (FISH) technique for detection of HER-2/neu gene amplification is also performed on breast cancer tissue removed at surgery, and measures HER-2/neu gene amplification present in cells. It directly tests DNA in the cancer cell to determine HER-2/neu status at the genetic level and reflects the number of HER-2/neu and 17 chromosome centromere FISH signals enumerated in 50 to 100 cells. Results are reported as a ratio of the number of HER-2 signals to 17 chromosome centromere signals. A ratio of less than 1.8 is within normal limits, a ratio of 1.8-2.0 is

Medical Policy: HER-2 TESTING IN INVASIVE BREAST OR GASTRIC CANCER USING FLUORESCENCE IN SITU HYBRIDIZATION (FISH) OR IMMUNOHISTOCHEMISTRY (IHC) ASSAYS

Policy Number: 2.02.31

Page: 3 of 8

equivocal and requires further testing, and a ratio of greater than 2 is consistent with amplification of HER-2/neu gene sequences. An example of a FISH HER-2/neu test is the PathVysion® test.

RATIONALE

The U. S. Food and Drug Administration (FDA) has given premarket approval for kits using the FISH technique for quantifying HER-2/neu gene amplification in formalin-fixed, paraffin-embedded breast cancer tissue specimens. These include but are not limited to the PathVysion® HER-2 DNA Probe Kit (Vysis, Inc.) approved in 2001, the INFORM® her-2/neu test kit (Ventana Medical Systems, Inc.) approved in 2000, and the DakoCytomation *Her2* FISH pharmDx™ kit (Denmark) approved in 2005. However, only the PathVysion® FISH test kit has been further indicated by the FDA to be an aid in the assessment of patients for whom Herceptin® (Trastuzumab) treatment is being considered. In April 2005 the FDA approved the Ariol HER-2/neu FISH software application (Applied Imaging Corp.) which allows the Ariol automated scanning microscope and image analysis system to detect amplification of the HER-2/neu gene via fluorescence in situ hybridization (FISH) in human breast cancer biopsy samples. The application and system are intended for use with a DNA probe kit (PathVysion HER-2/neu, Vysis, Inc.). The IHC HercepTest® (DAKO, Glostrup, Denmark) and the IHC Pathway® HER2 test (Ventana Medical Systems, Tucson, AZ) have FDA approval for determining the HER2 status of breast cancer tumors.

In 2013, the American Society of Clinical Oncology (ASCO) and the College of American Pathologists (CAP) published the updated results of an expert panel that conducted a systematic review of the literature and developed recommendations for optimal HER2 testing performance including new guidelines for consistent handling of samples. The guideline was reviewed by selected experts and approved by the board of directors for both organizations. The panel recommended that all newly diagnosed invasive breast cancers (primary site and/or metastatic site) be tested for HER2 status, ensuring that at least one tumor sample is tested for either HER2 protein expression (immunohistochemistry [IHC] assay) or (in situ hybridization [ISH assay]) for *HER2* gene amplification. The role of HER2-targeted therapy should be discussed when the HER2 test result is positive and if there is no apparent histopathologic discordance with HER2 testing. The decision to recommend HER2-targeted therapy should be delayed if the HER2 test result is equivocal. Mandatory retesting should be performed on the same specimen using the alternative test if the initial HER2 test result is equivocal or on an alternative specimen. If the HER2 test result is negative, HER2-targeted therapy should not be administered. If there is apparent histopathologic discordance with the HER2 test result, additional HER2 testing should be considered. A HER2 test result should be reported as indeterminate if technical issues prevent one or both tests (IHC and ISH) from being done in a tumor specimen, or prevent the test (or tests) from being reported as positive, negative, or equivocal. Confirm that the testing laboratory conforms to standards set for accreditation by CAP or an equivalent accreditation authority. In 2018, the American Society of Clinical Oncology (ASCO) and the College of American Pathologists (CAP) published the updated results from 2013 with two recommendations: first, immunohistochemistry (IHC) 2+ is defined as invasive breast cancer with weak to moderate complete membrane staining observed in greater than 10% of tumor cells. Second, if the initial HER2 test result in a core needle biopsy specimen of a primary breast cancer is negative, a new HER2 test may (not “must”) be ordered on the excision specimen based on specific clinical criteria.

A retrospective study evaluated the concordance between HER2 gene amplification determined by FISH and HER2 protein overexpression previously determined by IHC in breast cancer tissue specimens from women screened for three pivotal clinical trials, including one international trial, of trastuzumab (Herceptin®) at 54 centers. In this study, 5,998 breast cancer tissue specimens were divided into two groups: IHC score of 0/1+ and IHC score 2+/3+. 300 specimens from each group were randomly selected to determine HER2 amplification using the FISH assay. Assay agreement between FISH and specimens with IHC scores were 0 = 97%, 1+ = 93%, and 3+ = 89%. Only 24% of specimens with 2+ IHC showed HER2 amplification by FISH (76% disagreement in this IHC subgroup).

A randomized, controlled, multi-center clinical trial evaluated the predictive value of HER2 in a population of advanced breast cancer patients randomly treated either with single-agent doxorubicin or with single-agent docetaxel. Of 326 patients in the trial, tumor samples were available for 176 patients (54%). Different cohorts of patients identified by HER2 were examined to assess a possible relationship between HER2 status and treatment effect. In this trial, all positive IHC

Medical Policy: HER-2 TESTING IN INVASIVE BREAST OR GASTRIC CANCER USING FLUORESCENCE IN SITU HYBRIDIZATION (FISH) OR IMMUNOHISTOCHEMISTRY (IHC) ASSAYS

Policy Number: 2.02.31

Page: 4 of 8

cases received FISH to confirm HER2-positive status. HER2 positivity was observed in 20% of the study population. A statistically significant interaction was found between response rates to the study drugs and HER2 status.

Numerous studies with small sample sizes support that HER-2/neu status with an immunohistochemistry (IHC) score of 2+ should be confirmed with FISH testing, and that 3% to 7% of patients who are negative by IHC are found to be positive by FISH testing.

The 2020 National Comprehensive Cancer Network (NCCN) Practice Guidelines in Oncology for Breast Cancer recommend determination of HER2 status for all newly diagnosed or newly metastatic invasive breast cancers. The guidelines state that HER2 status can be assessed by measuring the number of HER2 gene copies (FISH assay) or by a complementary method in which the number of HER2 cell surface receptors is evaluated (IHC assay). The NCCN Breast Cancer Panel recommends selecting patients for trastuzumab (Herceptin) therapy who have tumors either positive for HER2 by FISH, or 3+ by IHC. Patients with tumors IHC 0, or 1+ for HER2, or FISH not amplified have very low rates of trastuzumab response and therefore therapy with trastuzumab is not warranted. Additionally either an IHC assay or a FISH assay can be used to make an initial assessment of HER2 tumor status, and all HER2 assays must be validated. Borderline IHC samples (e.g. IHC 2+) should be subjected to reflex testing by a validated complementary (e.g. FISH) method. Borderline FISH samples (e.g. an average HER2 gene/chromosome 17 ratio of 1.8-2.2 or an average HER2 gene copy number of greater than 4 to less than 6) should undergo: counting of additional cells, retesting by FISH, or reflex testing by a validated IHC method. A validated FDA-approved version of the FISH assay is recommended as the “gold standard” for confirmatory testing, when necessary. The NCCN panel endorses the use of College of American Pathologists protocols for reporting the pathological analysis of all breast specimens.

The 2020 National Comprehensive Cancer Network (NCCN) Practice Guidelines in Oncology for Gastric Cancer recommend determination of HER2 status for patients with inoperable, locally advanced, recurrent or metastatic adenocarcinoma of the stomach or EGJ for whom trastuzumab therapy is being considered. Assessment for tumor HER2 overexpression using immunohistochemistry (IHC) and fluorescence in situ hybridization (FISH) or other in situ hybridization method is recommended. The NCCN Guidelines panel recommends that HER2 immunohistochemistry (IHC) be ordered/performed first, followed by in situ hybridization methods in cases with 2+ (equivocal) expression by IHC. Positive (3+) or negative (0 to 1+) HER2 IHC results not require further ISH testing. Cases with HER2:CEP17 ration greater than or equal to 2 or an average HER2 copy number greater than or equal to 6 are considered positive by ISH/FISH.

CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member’s subscriber contract.*
- *CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.*
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*

CPT Codes

Code	Description
88341	Immunohistochemistry or immunocytochemistry, per specimen; each additional single antibody stain procedure (list separately in addition to code for primary procedure)
88344	Immunohistochemistry or immunocytochemistry, per specimen; each multiplex antibody stain procedure
88360	Morphometric analysis, tumor immunohistochemistry (e.g. HER-2/neu, estrogen receptor/progesterone receptor), quantitative or semiquantitative, per specimen, each single antibody stain procedure; manual

Medical Policy: HER-2 TESTING IN INVASIVE BREAST OR GASTRIC CANCER USING FLUORESCENCE IN SITU HYBRIDIZATION (FISH) OR IMMUNOHISTOCHEMISTRY (IHC) ASSAYS

Policy Number: 2.02.31

Page: 5 of 8

Code	Description
88361	Morphometric analysis, tumor immunohistochemistry (eg, Her-2/neu, estrogen receptor/progesterone receptor), quantitative or semiquantitative, per specimen, each single antibody stain procedure; using computer-assisted technology
88364	In situ hybridization (e.g., FISH), per specimen; each additional single probe stain procedure (list separately in addition to code for primary procedure)
88365	In situ hybridization (e.g. FISH), per specimen; initial single probe stain procedure
88367	Morphometric analysis, in situ hybridization, (quantitative or semi-quantitative); using computer-assisted technology, per specimen; initial single probe stain procedure
88368	Morphometric analysis, in situ hybridization, (quantitative or semi-quantitative), manual, per specimen; initial single probe stain procedure
88369	Morphometric analysis, in situ hybridization, (quantitative or semi-quantitative), manual, per specimen; each additional single probe stain procedure (list separately in addition to code for primary procedure)
88373	Morphometric analysis, in situ hybridization, (quantitative or semi-quantitative), using computer-assisted technology, per specimen; each additional single probe stain procedure (list separately in addition to code for primary procedure)
88374	Morphometric analysis, in situ hybridization, (quantitative or semi-quantitative), using computer-assisted technology, per specimen; each multiplex probe stain procedure
88377	Morphometric analysis, in situ hybridization, (quantitative or semi-quantitative), manual, per specimen; each multiplex probe stain procedure
0009U	Oncology (breast cancer), ERBB2 (HER2) copy number by FISH, tumor cells from formalin fixed paraffin embedded tissue isolated using image-based dielectrophoresis (DEP) sorting, reported as ERBB2 gene amplified or non-amplified (DEPArray™ HER2, PacificDx)

Copyright © 2020 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
No code(s)	

ICD10 Codes

Code	Description
C16.0-C16.9	Malignant neoplasm of stomach (code range)
C50.0-C50.9	Malignant neoplasm of breast (code range)
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
C78.80	Secondary malignant neoplasm of unspecified digestive organ
C78.89	Secondary malignant neoplasm of other digestive organs
D00.2	Carcinoma in situ of stomach
D37.1-D37.5	Neoplasm of uncertain behavior of digestive organs (code range)

Medical Policy: HER-2 TESTING IN INVASIVE BREAST OR GASTRIC CANCER USING FLUORESCENCE IN SITU HYBRIDIZATION (FISH) OR IMMUNOHISTOCHEMISTRY (IHC) ASSAYS

Policy Number: 2.02.31

Page: 6 of 8

Code	Description
Z85.3	Personal history of malignant neoplasm of breast

REFERENCES

- *Bagaria SP, et al. Personalizing breast cancer staging by the inclusion of ER, PR, and HER2. JAMA Surg 2014 Feb;149(2):125-9.
- Barsouk A, et al. Targeted therapies and immunotherapies in the treatment of esophageal cancers. Med. Sci. 2019;7(10):100.
- *Bartlett JM, et al. External quality assurance of HER2 FISH and ISH testing: Three years of the UK National External Quality Assurance Scheme. Am J Clin Pathol 2009;131:106-11.
- *Chao WR, et al. Assessing the HER2 status in mucinous epithelial ovarian cancer on the basis of the 2013 ASCO/CAP guideline update. Am J Surg Pathol 2014 Sep;38(9):1227-34.
- Dijksterhuis WPM, et al. Increased assessment of HER2 in metastatic gastroesophageal cancer patients: a nationwide population-based cohort study. Gastric Cancer 2020 Jan 11 [Epub ahead of print].
- *Chua TC, et al. Clinicopathologic factors associated with HER2-positive gastric cancer and its impact on survival outcomes – a systematic review. Int J Cancer 2012 Jun 15;130(12):2845-56.
- *Elimova E, et al. Medical management of gastric cancer: a 2014 update. World J Gastroenterol 2014 Oct 14;20(38):13637-47.
- *Engelstaedter V, et al. Her-2/neu and topoisomerase I α in advanced breast cancer: a comprehensive FISH analysis of 245 cases. Diagn Mol Pathol 2012 Jun;21(2):77-83.
- Feng Y, et al. HER2 as a potential biomarker guiding adjuvant chemotherapy in stage II colorectal cancer. Eur J Surg Oncol. 2019 Feb;45(2):167-173.
- *Fisher SB, et al. HER2 in resected gastric cancer: is there prognostic value? J Surg Oncol 2014 Feb;109(2):61-6.
- Guo L, et al. Analysis of molecular subtypes for the increased HER2 equivocal cases caused by application of the updated 2013 ASCO/CAP HER2 testing guidelines in breast cancer. Breast Cancer Res Treat. 2017 Nov;166(1):77-84.
- Jeong JH, et al. HER2 amplification and cetuximab efficacy in patients with metastatic colorectal cancer harboring wild-type RAS and BRAF. Clin Colorectal Cancer 2017 Sep;16(3):e147-e152.
- Kim EK, et al. The frequency and clinical impact of HER2 alterations in lung adenocarcinoma. PLoS One 2017 Feb 1;12(2):e0171280.
- Lokich E, et al. Molecular markers in uterine serous cancer: correlation between endometrial biopsy and hysterectomy specimens. Gynecol Oncol Rep. 2019 Apr 25;29:98-101.
- Meric-Bernstam F, et al. Advances in HER2-targeted therapy: novel agents and opportunities beyond breast and gastric cancer. Clin Cancer Res 2019 Apr 1;25(7):2033-2041.
- Meng X, et al. Expression of human epidermal growth factor receptor-2 in resected rectal cancer. Medicine (Baltimore) 2015 Nov;94(47):e2108.
- *Middleton LP, et al. Implementation of American Society of Clinical Oncology/College of American Pathologists HER2 guideline recommendations in a tertiary care facility increases HER2 immunohistochemistry and Fluorescence In Situ Hybridization concordance and decreases the number of inconclusive cases. Arch Pathol Lab Med 2009;133:775–80.
- Murray NP, et al. Possible role of HER-2 in the progression of prostate cancer from primary tumor to androgen independence. Asian Pac J Cancer Prev 2015;16(15):6615-9.

Medical Policy: HER-2 TESTING IN INVASIVE BREAST OR GASTRIC CANCER USING FLUORESCENCE IN SITU HYBRIDIZATION (FISH) OR IMMUNOHISTOCHEMISTRY (IHC) ASSAYS

Policy Number: 2.02.31

Page: 7 of 8

National Comprehensive Cancer Network, Inc. Practice guidelines in oncology Breast Cancer - version 2. 2020 [http://www.nccn.org/professionals/physician_gls/pdf/breast.pdf] accessed 2/09/20.

National Comprehensive Cancer Network, Inc. Practice guidelines in oncology Colon Cancer - version 1. 2020 [https://www.nccn.org/professionals/physician_gls/pdf/colon_blocks.pdf] accessed 2/09/20.

National Comprehensive Cancer Network, Inc. Practice guidelines in oncology Gastric Cancer - version 4. 2019 [http://www.nccn.org/professionals/physician_gls/pdf/gastric.pdf] accessed 2/09/20.

National Comprehensive Cancer Network, Inc. Practice guidelines in oncology NonSmall Lung Cancer - version 2. 2020 [https://www.nccn.org/professionals/physician_gls/pdf/nscl_blocks.pdf] accessed 02/09/20.

Nedjadi T, et al. Prognostic value of HER2 status in bladder transitional cell carcinoma revealed by both IHC and BDISH. BMC Cancer 2016 Aug 19;16:653.

*Park MM, et al. ER and PR immunohistochemistry and HER2FISH versus Oncotype DX: implications for breast cancer treatment. Breast J 2014 Jan-Feb;20(1):37-45.

*Perez EA, et al. *HER2* and Chromosome 17 effect on patient outcome in the N9831 adjuvant Trastuzumab trial. J Clin Oncol 2010 Oct 1;28(28):4307-15.

Prieto M, et al. Long term recurrence free survival in a stage IV gallbladder cancer treated with chemotherapy plus trastuzumab and salvage liver resection. Ann Hepatobiliary Pancreat Surg 2019;23:403-407.

*Schink JC, et al. Biomarker testing for breast, lung, and gastroesophageal cancers at NCI designated cancer centers. J Natl Cancer Inst 2014 Sep 12;106(10).

*Seo AN, et al. HER2 status in colorectal cancer: its clinical significance and the relationship between HER2 gene amplification and expression. PLoS One 2014 May 30;9(5):e98528.

Takegawa N and Yonesaka K. HER2 as an emerging oncotarget for colorectal cancer treatment after failure of anti-epidermal growth factor receptor therapy. Clin Colorectal Cancer 2017 Dec;16(4):247-251.

Teplinsky E, et al. Targeting HER2 in ovarian and uterine cancers: challenges and future directions. Gynecol Oncol 2014 Nov;135(2):364-70.

Van Cutsem E, et al. HER2 screening data from ToGA: targeting HER2 in gastric and gastroesophageal junction cancer. Gastric Cancer 2015 Jul;18(3):476-84.

*Wagner AD et al. Chemotherapy for advanced gastric cancer. Cochrane Database Syst Rev 2010 Mar 17;(3):CD004064.

Wolff AC, et al. Receptor 2 Testing in Breast Cancer: American Society of Clinical Oncology/College of American Pathologists Clinical Practice Guideline Update. J Clin Oncol. 2013 Nov 1;31(31):3997-4013.

Wolff AC, et al. Human Epidermal Growth Factor Receptor 2 Testing in Breast Cancer: American Society of Clinical Oncology/College of American Pathologists Clinical Practice Guideline Focused Update. Arch Pathol Lab Med 2018 Nov;142:1364-1382.

Wu SW, et al. Does overexpression of HER-2 correlate with clinicopathological characteristics and prognosis in colorectal cancer? Evidence from a meta-analysis. Diagn Pathol 2015 Aug 16;10:144.

Xu FP, et al. Impact of repeat HER2 testing after initial equivocal HER@ FISH results using 2013 ASCO/CAP guidelines. Breast Cancer Res Treat. 2017 Dec;166(3):757-764.

Xu QQ, et al. HER2 amplification level is not a prognostic factor for HER2-Positive breast cancer with trastuzumab-based adjuvant treatment: a systematic review and meta-analysis. Oncotarget 2016 Sep27;7(30):63571-63582.

Zare S, et al. Implementation of the 2018 American Society of Clinical Oncology/College of American Pathologists Guidelines on HER2/neu assessment by FISH breast cancers: predicted impact in a single institutional cohort. Mod Pathol 2019 Nov;32(11):1566-1573.

Medical Policy: HER-2 TESTING IN INVASIVE BREAST OR GASTRIC CANCER USING FLUORESCENCE IN SITU HYBRIDIZATION (FISH) OR IMMUNOHISTOCHEMISTRY (IHC) ASSAYS

Policy Number: 2.02.31

Page: 8 of 8

*Zhang H, et al. HER-2 gene amplification by fluorescence in situ hybridization (FISH) compared with immunohistochemistry (IHC) in breast cancer: a study of 528 equivocal cases. Breast Cancer Res Treat 2012 Jul;134(2):743-9.

*Key Article

KEY WORDS

FISH, HercepTest, HER-2 overexpression, HER-2 amplification, IHC, immunohistochemistry, PathVysion

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently no National Coverage Determination (NCD) or Local Coverage Determination (LCD) for HER-2 testing in invasive breast cancer using Fluorescence in situ hybridization (FISH) or immunohistochemistry IHC) assays.