



MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	POSITRON EMISSION TOMOGRAPHY (PET) NON-ONCOLOGIC APPLICATIONS
Policy Number	6.01.07
Category	Technology Assessment
Effective Date	10/18/01
Revised Date	10/18/01, 01/17/02, 10/16/02, 01/16/03, 10/15/03, 10/20/04, 10/20/05, 11/16/06, 08/16/07, 08/21/08, 09/17/09, 12/16/10, 01/20/11, 12/15/11, 01/17/13, 05/22/14, 02/19/15, 02/18/16, 02/16/17, 02/15/18, 06/20/19, 03/19/20
Product Disclaimer	<ul style="list-style-type: none"> • If a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan product) or a Medicaid product covers a specific service, medical policy criteria apply to the benefit. • If a Medicare product covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit.

POLICY STATEMENT

- I. Based upon our criteria and assessment of peer reviewed literature, FDG positron emission tomography (PET) using a full ring dedicated PET scanner is considered **medically appropriate** for the following indications:
- A. Epileptic Seizures:
 1. Seizure disorders with failed response to medical therapy when being considered for resection of suspected epileptogenic focus in a region of the brain accessible by surgery.
 - B. To differentiate Alzheimer's disease (AD) from frontotemporal lobe dementia (FTLD) in patients with a recent diagnosis of dementia, when **ALL** of the following are present:
 1. Patient meets diagnostic criteria for AD and FTLD; and
 2. Patient has a documented cognitive decline of at least six months' duration; and
 3. Evaluation has ruled out specific alternative neurodegenerative diseases or causative factors; and
 4. Cause of clinical symptoms is uncertain; and
 5. Results are expected to help clarify the diagnosis between FTLD and AD and help guide future treatment.
 - C. To evaluate patients suspected of having encephalitis, including autoimmune encephalitis, if diagnosis remains unclear after evaluation with brain MRI, CSF analysis, and lab testing, including serology, if appropriate.
- II. Based upon our criteria and assessment of the peer-reviewed literature, the use of beta amyloid PET imaging using amyloid specific tracers (e.g., Amyvid™, Vizamy™, Neuraceq™) for dementia has not been medically proven to be effective and, therefore, is considered **investigational**.
- III. Based upon our criteria and assessment of the peer-reviewed literature, the use of PET scanning has not been medically proven to be effective and, therefore, is considered **investigational** for all other indications, including, but not limited to:
- A. Anorexia Nervosa;
 - B. Auto-immune disorders with CNS manifestations, including Behcets' syndrome and lupus erythematosus;
 - D. Cerebral blood flow in newborns;
 - E. Cerebrovascular diseases, including arterial occlusive disease (arteriosclerosis, atherosclerosis), carotid artery disease, cerebral aneurysm, cerebrovascular malformations (AVM) hemorrhage, infarct, and ischemia;
 - F. Chronic fatigue syndrome;

Medical Policy: POSITRON EMISSION TOMOGRAPHY (PET) NON-ONCOLOGIC APPLICATIONS

Policy Number: 6.01.07

Page: 2 of 7

- G. Degenerative motor neuron diseases, including amyotrophic lateral sclerosis (ALS), Friedreich's ataxia, olivopontocerebellar atrophy, Parkinson's disease, progressive supranuclear palsy, Shy-Drager syndrome, spinocerebellar degeneration, Steele-Richardson-Olszewski disease, and Tourette's syndrome;
- H. Dementias, including, dementia with Lewy-bodies, multi-infarct dementia, Pick's disease, presenile dementia, Alzheimer's disease, and frontotemporal dementia, except as listed in Policy Statement I.E.;
- I. Demyelinating diseases, such as multiple sclerosis;
- J. Developmental, congenital, or inherited disorders, including adrenoleukodystrophy, Down's syndrome, Kinky-hair disease (Menkes' syndrome), Sturge-Weber syndrome (encephalofacial angiomatosis), and the phakomatoses;
- J. Diagnosis and non-surgical treatment of epilepsy and convulsive disorders;
- K. Fever of unknown origin, infectious process;
- L. Giant cell arteritis;
- M. Inflammatory bowel disease;
- N. Inflammation of unknown origin;
- O. Joint replacement follow-up;
- P. Migraines;
- Q. Mycobacterium infection;
- R. Nutritional or metabolic diseases and disorders, including acanthocytes, hepatic encephalopathy, hepatolenticular degeneration, metachromatic leukodystrophy, mitochondrial disease, and subacute necrotizing encephalomyelopathy;
- S. Post-traumatic stress disorder;
- T. Psychiatric disease and disorders, including affective disorders, depression, obsessive-compulsive disorder, psychomotor disorders, schizophrenia;
- U. Pulmonary diseases, including adult respiratory distress syndrome, diffuse panbronchiolitis, emphysema, obstructive lung disease, and pneumonia;
- V. Pyogenic infections, including aspergillosis and encephalitis;
- W. Sarcoidosis (cardiac sarcoid - please refer to *Corporate Medical Policy #6.01.41 Positron Emission Tomography (PET) Cardiac Applications*);
- X. Sick building syndrome;
- Y. Spondylodiscitis;
- Z. Substance abuse, including CNS effects of alcohol, cocaine, and heroin;
- AA. Trauma, including brain injury and carbon monoxide poisoning;
- BB. Vasculitis;
- CC. Viral infections, including acquired immune deficiency syndrome (AIDS), AIDS dementia complex, Creutzfeldt-Jakob syndrome, progressive multifocal leukoencephalopathy, progressive rubella encephalopathy, and subacute sclerosing panencephalitis.

Refer to Corporate Medical Policy #6.01.29 Positron Emission Tomography-Oncologic Applications.

Refer to Corporate Medical Policy #6.01.41 Positron Emission Tomography (PET) Cardiac Application

Refer to Corporate Medical Policy #11.01.03 Experimental or Investigational Services.

POLICY GUIDELINES

The Federal Employees Health Benefit Program (FEHBP/FEP) requires that procedures, devices or laboratory tests approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational and, thus, these procedures, devices or laboratory tests may be assessed only on the basis of their medical necessity.

Medical Policy: POSITRON EMISSION TOMOGRAPHY (PET) NON-ONCOLOGIC APPLICATIONS

Policy Number: 6.01.07

Page: 3 of 7

DESCRIPTION

Positron emission tomography (PET) is an imaging technology that can reveal metabolic information in various tissue sites. The metabolic information is what distinguishes it from other imaging modalities such as magnetic resonance imaging (MRI) and computed tomography (CT), which provide primarily anatomic information. PET scans measure concentrations of radioactive chemicals that are partially metabolized in the body and are based on the use of positron emitting radionuclide tracers coupled to organic molecules such as glucose, ammonia, or water. Dedicated PET scanners consist of multiple detectors arranged in a full or partial ring around the patient.

A variety of radiotracers are used for PET scanning, including fluorine-18, rubidium-82, ammonia N-13, carbon-11, oxygen-15, and nitrogen-13. Fluorine-18 is often coupled with fluoro-deoxyglucose (FDG) as a means of detecting glucose metabolism, which, in turn, reflects the metabolic activity, and, thus, viability, of the target tissue. Because of their short half-life, tracers must be made locally. With the exception of fluorine and rubidium, all the tracers must be manufactured with an on-site cyclotron.

Florbetapir (Amyvid™, Avid Radiopharmaceuticals), a radioactive dye for visualization of amyloid plaque in the brain, was approved by the FDA in 2012. The FDA document prepared for the advisory committee meeting indicated that, while florbetapir may detect pathology, there could be no claim of disease detection, as beta amyloid aggregates can be found in cognitively normal elderly individuals, as well as patients with Alzheimer's disease (AD). Amyvid™ is indicated for PET imaging of the brain, to estimate beta-amyloid neuritic plaque density in adult patients with cognitive impairment who are being evaluated for AD and other causes of cognitive decline. A second radioactive dye, Flutemetamol F18 injection (Vizamyl™, GE Healthcare), was approved by the FDA in October, 2013. Flutemetamol F18 is not indicated to predict the development of AD or to check how patients respond to treatment for AD. Flutemetamol F18 PET images should be interpreted only by health care professionals who successfully complete training in an image interpretation program. In March 2014, the FDA approved a third radioactive dye, florbetaben F18 (Neuraceq™; Piramal Life Sciences, Matran, Switzerland).

RATIONALE

The FDA has approved the scanner and imaging hardware for PET as being substantially equivalent to x-ray CT. The FDA requires PET radiotracers to be approved through a new drug approval (NDA) process. Because PET radiotracers have an extremely short half-life, they must be produced in the clinical setting. The FDA also regulates drug manufacturing processes in PET facilities. In 1991, the FDA approved the use of Rubidium 82 (Rb 82) as a myocardial perfusion tracer and, in 1999, approved the use of ammonia N-13 as a myocardial perfusion tracer.

Clinical evidence supports that the use of Rubidium 82 (Rb-82) PET and ammonia N-13 PET scans in clinical practice has the potential to improve net health outcomes through changes in patient management. Studies demonstrate that both tracers have high reliability and validity in the evaluation of myocardial perfusion.

Clinical evidence is inadequate to support the use of FDG PET for routine use in the diagnostic evaluation of dementia. Although FDG PET scanning appears to have promise for use as an adjunct to clinical diagnosis of Alzheimer's disease, further prospective studies are needed to establish the value that it brings to diagnosis over and above a competent clinical diagnosis. A National Institute on Aging longitudinal, five-year, prospective trial, the Alzheimer's Disease Neuroimaging Initiative (ADNI), plans to include 800 participants aged 55-90 years (400 with mild cognitive impairment, 200 with Alzheimer's disease, 200 normal participants) to be followed for two years. At 58 sites in the U.S. and Canada ADNI will compare neuroimaging (PET and MRI), biological, and clinical information. It will seek correlations among data that will track the progression of memory loss from its earliest stages, and identify critical markers that response to treatments aimed at slowing progression of mild cognitive impairment and Alzheimer's disease. Enrollment began in early 2006 and the end date is anticipated to be October 2009.

A 2013 BlueCross BlueShield Technical Assessment concluded that beta amyloid imaging with PET to evaluate suspected AD and other causes of cognitive decline does not meet the TEC criteria, based on the lack of direct evidence for clinical utility. The test is not likely to be useful for confirming AD in patients who present with cognitive impairment. It may have a role in ruling out AD, but this has yet to be established with certainty. Questions also remain

Medical Policy: POSITRON EMISSION TOMOGRAPHY (PET) NON-ONCOLOGIC APPLICATIONS

Policy Number: 6.01.07

Page: 4 of 7

about the use of this test outside of the investigational setting, particularly regarding the accuracy of visual interpretation of images and how best to apply this test in routine clinical practice.

Clinical evidence in the form of small prospective and retrospective studies totaling 166 patients, and a meta-analysis of 19 studies, support that FDG PET is highly accurate in diagnosing chronic osteomyelitis.

Several studies with methodologic flaws indicate that there are instances in which PET may be helpful in the diagnosis of fever of unknown origin and infection. However, clinical evidence is not sufficient to consider these indications medically appropriate.

FDG-PET has been investigated for potential use in the diagnosis and follow-up of giant cell arteritis. Clinical evidence consists of small case series, retrospective studies, and case reports. Although some reports consider PET promising for this indication, results need to be confirmed in larger prospective studies. The limited spatial resolution of PET scanners is a technical limitation that prevents the detection of metabolic signals within anatomical structures smaller than four to five mm in size. In addition, the physiological uptake of FDG by the grey matter of the brain obscures FDG uptake within the temporal arteries.

CODES

- *Eligibility for reimbursement is based upon the benefits set forth in the member's subscriber contract.*
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- *Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.*

CPT Codes

Code	Description
78608	Brain imaging, positron emission tomography, (PET); metabolic evaluation
78609	perfusion evaluation
78811	Positron emission tomography (PET) imaging; limited area (e.g., chest, head/neck)
78812	skull base to mid-thigh
78813	whole body
78814	Positron emission tomography (PET) with concurrently acquired computed tomography (CT) for attenuation correction and anatomical localization imaging; limited area (e.g. chest, head/neck)
78815	skull base to mid-thigh
78816	whole body

Copyright © 2020 American Medical Association, Chicago, IL

HCPCS Codes

Code	Description
A9526	Nitrogen N-13 ammonia, diagnostic, per study dose, up to 40 millicuries
A9552	Fluorodeoxyglucose F-18 FDG, diagnostic, per study dose, up to 45 millicuries
A9555	Rubidium Rb-82, diagnostic, per study dose, up to 60 millicuries
A9586 (E/I)	Florbetapir F18, diagnostic, per study dose, up to 10 millicuries
A9598 (E/I)	Positron emission tomography radiopharmaceutical, diagnostic, for non-tumor identification, not otherwise classified

Medical Policy: POSITRON EMISSION TOMOGRAPHY (PET) NON-ONCOLOGIC APPLICATIONS

Policy Number: 6.01.07

Page: 5 of 7

Code	Description
Q9982 (E/I)	Flutemetamol F18, diagnostic, per study dose, up to 5 millicuries
Q9983 (E/I)	Florbetaben F18, diagnostic, per study dose, up to 8.1 millicuries
S8085 (E/I)	Fluorine-18 fluorodeoxyglucose (F-18 FDG) imaging using dual-head coincidence detection system (non-dedicated PET scan)

ICD10 Codes

Code	Description
D33.0-D33.2	Benign neoplasm of brain and other parts of central nervous system (code range)
D43.0-D43.4	Neoplasm of uncertain behavior of brain and spinal cord (code range)
D49.6	Neoplasm of unspecified behavior of brain
G40.001- G40.219	Localization-related (focal) (partial) idiopathic epilepsy and epileptic syndromes with seizures of localized onset (code range)
G40.301- G40.3119	Generalized idiopathic epilepsy and epileptic syndromes (code range)
G40.901- G40.919	Epilepsy, unspecified (code range)

REFERENCES

BlueCross BlueShield Association. Beta amyloid imaging with positron emission tomography (PET) for Alzheimer's Disease. Medical Policy Reference Manual. Policy #6.01.55. 2019 Sep 12.

*BlueCross BlueShield Association. FDG using camera-based imaging (FDG-SPECT). Medical Policy Reference Manual. Policy #6.01.27. Archived 2009 Dec 03.

BlueCross BlueShield Association. Miscellaneous applications of positron emission tomography (PET). Medical Policy Reference Manual. Policy #6.01.06. 2019 Sep 12.

Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Beta amyloid imaging with positron emission tomography (PET) for evaluation of suspected Alzheimer's Disease or other causes of cognitive decline. TEC Assessments 2013;27:5.

*Castaigne C, et al. Clinical value of FDG-PET/CT for the diagnosis of human immunodeficiency virus-associated fever of unknown origin: a retrospective study. Nuclear Medicine Communic 2009;30:41-47.

Chet lat G, et al. Amyloid imaging in cognitively normal individuals, at-risk populations and preclinical Alzheimer's disease. Neuroimage Clin 2013 Mar 5;2:356-65.

*deWinter F, et al. Fluorine-18-fluorodeoxyglucose-positron emission tomography: a highly accurate imaging modality for the diagnosis of chronic musculoskeletal infections. J Bone Joint Surg Am 2001 May;83-A(5):651-660.

Doraiswamy PM, et al. Florbetapir F 18 amyloid Pet and 36-month cognitive decline: a prospective multicenter study. Mol Psychiatry 2014 Sep;19(9):1044-51.

*Duke Evidence-based Practice Center. Use of positron emission tomography and other neuroimaging techniques in the diagnosis and management of Alzheimer's disease and dementia. Technology Assessment prepared for the Agency for HealthCare Research and Quality. Contract No. 290-97-0014, Task Order 7. 2001 Dec 14.

*Gill SS, et al. The value of positron emission tomography in the clinical evaluation of dementia. J Am Geriatr Soc 2003 Feb;51(2):258-264.

Medical Policy: POSITRON EMISSION TOMOGRAPHY (PET) NON-ONCOLOGIC APPLICATIONS

Policy Number: 6.01.07

Page: 6 of 7

*Guhlmann A, et al. Fluorine-18-FDG PET and technetium-99m antiglycyl antibody scintigraphy in chronic osteomyelitis. J Nucl Med 1998 Dec;39(12):2145-2152.

*Hartmann A, et al. Diagnostic value of 18F-FDG PET/CT in trauma patients with suspected chronic osteomyelitis. Eur J Nucl Med Mol Imaging 2007 May;34(5):704-714.

*Hove JD, et al. Simultaneous cardiac output and regional myocardial perfusion determination with PET and nitrogen 13 ammonia. J Nucl Cardiol 2003 Jan-Feb;10(1):28-33.

*Knopman DS, et al. Practice parameter: Diagnosis of dementia (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. Neurol 2001;56:1143-1153.

*Kwee TC, et al. FDG-PET for diagnosing prosthetic joint infection: a systematic review and metaanalysis. Eur J Nucl Med Mol Imaging 2008;35:2122–2132.

Lim YY, et al. Effect of amyloid on memory and non-memory decline from preclinical to clinical Alzheimer's disease. Brain 2014 Jan;137(1):221-31.

Ma Y, et al. Predictive accuracy of amyloid imaging for progression from mild cognitive impairment to Alzheimer disease with different lengths of follow-up. Medicine (Baltimore) 2014 Dec;93(27):e150.

Mattsson N, et al. Independent information from cerebrospinal fluid amyloid- β and florbetapir imaging in Alzheimer's disease. Brain 2015 Mar;138(Pt 3):772-83.

Nazar AH, et al. Spectrum of 18F-FDG PET/CT findings in patients presenting with fever of unknown origin. AJR Am J Roentgenol 2012 Jul;199(1):175-85.

*Panegyres PK, et al. Fluorodeoxyglucose–Positron Emission Tomography in the differential diagnosis of early-onset dementia: a prospective, community-based study. BMC Neurol 2009;9:41.

Rabinovici , et al. Testing and disclosures related to amyloid imaging and Alzheimer's disease: Common questions and fact sheet summary. Alzheimers Dement. 2016 Apr;12(4):510-5.

*Silverman DH, et al. Positron emission tomography in evaluation of dementia: regional brain metabolism and long-term outcome. JAMA 2001 Nov 7;286(17):2120-2127.

Torosyan N, et al. Neuronuclear imaging in the evaluation of dementia and mild decline in cognition. Semin Nucl Med 2012 Nov;42(6):415-422.

Witte MM, et al. Association between clinical measures and florbetapir F18PET neuroimaging in mild or moderate Alzheimer's disease dementia. J Neuropsychiatry Clin Neurosci 2014 Summer;26(3):214-20.

*Yuan Y, et al. Fluorodeoxyglucose–Positron-Emission Tomography, Single-Photon Emission Tomography, and structural MR imaging for prediction of rapid conversion to Alzheimer Disease in patients with mild cognitive impairment: a meta-analysis. AJNR Am J Neuroradiol 2009;30(2):404-410.

*Key Article

KEY WORDS

FDG PET, FDG SPECT, Gamma Camera, Ammonia N-13, Rubidium 82.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently a National Coverage Determination (NCD) for PET scans. Please refer to the following NCD website for Medicare Members: <http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=211&ncdver=4&bc=AgAAgAAAAAA&>

There is currently a National Coverage Determination (NCD) for FDG PET for Dementia and Neurodegenerative Diseases. Please refer to the following NCD website for Medicare Members: <http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=288&ncdver=3&bc=BAABAAAAAA&>

Medical Policy: POSITRON EMISSION TOMOGRAPHY (PET) NON-ONCOLOGIC APPLICATIONS

Policy Number: 6.01.07

Page: 7 of 7

There is currently a National Coverage Determination (NCD) for Beta Amyloid Positron Tomography in Dementia and Neurodegenerative Disease. Please refer to the following NCD website for Medicare Members:

<http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=356&ncdver=1&bc=AgAAgAAAAAAAAA%3d%3d&>