

MEDICAL POLICY

MEDICAL POLICY DETAILS	
Medical Policy Title	Spinal Cord Stimulation/Dorsal Column Stimulation
Policy Number	7.01.51
Category	Technology Assessment
Original Effective Date	11/15/01
Committee Approval Date	09/19/02, 09/18/03, 07/15/04, 07/21/05, 05/18/06, 04/19/07, 06/19/08, 05/28/09, 04/22/10, 03/17/11, 03/15/12, 06/19/14, 08/20/15, 10/20/16, 10/19/17, 06/21/18, 12/20/18, 06/20/19, 08/20/20, 04/15/21, 09/16/21, 05/19/22, 05/18/23, 10/17/24
Current Effective Date	02/01/25
Archived Date	(03/21/13-06/19/14)
Archive Review Date	N/A
Product Disclaimer	<ul style="list-style-type: none"> • Services are contract dependent; if a product excludes coverage for a service, it is not covered, and medical policy criteria do not apply. • If a commercial product (including an Essential Plan or Child Health Plus product), medical policy criteria apply to the benefit. • If a Medicaid product covers a specific service, and there are no New York State Medicaid guidelines (eMedNY) criteria, medical policy criteria apply to the benefit. • If a Medicare product (including Medicare HMO-Dual Special Needs Program (DSNP) product) covers a specific service, and there is no national or local Medicare coverage decision for the service, medical policy criteria apply to the benefit. • If a Medicare HMO-Dual Special Needs Program (DSNP) product DOES NOT cover a specific service, please refer to the Medicaid Product coverage line.

POLICY STATEMENT

- I. Based upon our criteria and assessment of the peer-reviewed literature, implantation of a spinal cord stimulator (SCS) (non-high-frequency or high-frequency) has been medically proven to be effective and, therefore, is considered **medically appropriate** for treatment of patients with failed back surgery syndrome (FBSS) with intractable neuropathic leg pain, as follows:
- A. A short-term trial (e.g., greater than 48 hours) of spinal cord stimulation (non-high-frequency or high-frequency [HF 10 SCS]), when **ALL** of the following criteria are met:
1. Patient has failed at least six (6) consecutive months of conservative treatment that includes **all** of the following:
 - a. exercise;
 - b. manual therapy;
 - c. patient education;
 - d. psychosocial support/cognitive behavioral therapy; **and**
 - e. medications to include nonsteroidal anti-inflammatory drugs [NSAIDS] or analgesics;
 2. Surgical intervention is not indicated, or the patient does not wish to proceed with spinal surgery; **and**
 3. An attestation by a behavioral health provider (i.e., a face-to-face or virtual assessment, with or without psychological questionnaires) reveals no evidence of an inadequately controlled behavioral health condition/issue (e.g., substance use disorder, depression, or psychosis) that would impact perception of pain, and/or negatively impact the success of an SCS or contraindicate its placement. (See Policy Guidelines).
- B. Permanent implantation of an SCS (non-high-frequency or high-frequency [HF 10 SCS]) when at least a 50% reduction in pain has been demonstrated during a short-term trial of spinal cord stimulation (SCS).

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 2 of 16

- II. Based upon our criteria and assessment of the peer-reviewed literature, use of a non-high-frequency dorsal column SCS has been medically proven to be effective and, therefore, is considered **medically appropriate** for the treatment of chronic, intractable pain secondary to complex regional pain syndrome (CRPS)/reflex sympathetic dystrophy (RSD) only of the upper and lower extremities, as follows:
- A. A short-term trial (e.g., greater than 48 hours) of a non-high-frequency SCS, when **ALL** the following criteria are met:
1. Patient's diagnosis of CRPS/RSD, as evidenced by **all** the following:
 - a. Patient has continuing pain that is disproportionate to any inciting event;
 - b. Patient reports **at least one (1)** of the symptoms in **three (3) out of the four (4)** following categories:
 - i. Sensory: hyperesthesia;
 - ii. Vasomotor: temperature asymmetry, skin color changes, or skin color asymmetry;
 - iii. Sudomotor/edema: edema, sweating changes, or sweating asymmetry; **and/or**
 - iv. Motor/trophic: decreased range of motion, motor dysfunction (weakness, tremor, dystonia), trophic changes (hair, nail, skin);
 - c. On physical examination, patient displays **at least one (1)** of the signs in **two (2) or more** of the following categories:
 - i. Sensory: evidence of hyperalgesia (to pinprick) or allodynia (to light touch),
 - ii. Vasomotor: evidence of temperature asymmetry, skin color changes, or asymmetry,
 - iii. Sudomotor/edema: evidence of edema, sweating changes, or sweating asymmetry, **and/or**
 - iv. Motor/trophic: evidence of decreased range of motion, motor dysfunction (weakness, tremor, dystonia) or trophic changes (hair, nail, skin); **and**
 - d. There is/are no other medical or psychological diagnoses and/or results of relevant studies (e.g., imaging, electrodiagnostic testing, laboratory testing, etc.) that are concordant with the presenting signs and symptoms;
 2. Limited to only the extremities and not to the head/face/neck, trunk, perineum/pelvis, or abdominal viscera;
 3. Patient has failed at least six (6) consecutive months of conservative treatment that includes **all** of the following:
 - a. exercise;
 - b. manual therapy;
 - c. patient education;
 - d. psychosocial support/cognitive behavioral therapy; **and**
 - e. medications to include nonsteroidal anti-inflammatory drugs [NSAIDS] or analgesics;
 4. Surgical intervention is not indicated; **and**
 5. An attestation by a behavioral health provider (i.e., a face-to-face or virtual assessment, with or without psychological questionnaires) reveals no evidence of an inadequately controlled behavioral health condition/issue (e.g., substance use disorder, depression, or psychosis) that would impact perception of pain and/or negatively impact the success of an SCS or contraindicate its placement.
- B. Permanent implantation of a non-high-frequency dorsal column SCS, when at least a 50% reduction in pain has been demonstrated during a short-term trial of spinal cord stimulation.
- III. Based upon our criteria and assessment of the peer-reviewed literature, use of a non-high-frequency dorsal column SCS has been medically proven to be effective and, therefore, is considered **medically appropriate** for treatment of patients with chronic, intractable pain secondary to chronic critical limb ischemia (CLI), as follows:
- A. A short-term trial (e.g., greater than 48 hours) of a non-high-frequency dorsal column SCS, when **ALL** the following criteria are met:
1. Attestation from a vascular surgeon that the individual is not a suitable candidate for vascular reconstruction;
 2. Patient has a diagnosis of CLI when **all** of the following criteria are met:
 - a. Rutherford Classification Grade II, Category 4 (see Description section below), ischemic limb rest pain that is characterized by **both** of the following:

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 3 of 16

- i. resting ankle pressure less than 40 mmHg, flat or barely pulsatile ankle or metatarsal pulse volume recording; **and**
 - ii. toe pressure less than 30 mmHg;
 3. Advanced imaging (i.e., angiographic imaging, computed tomography [CT] scan or magnetic resonance imaging [MRI]) demonstrates multi-level disease with absence of named vessel with flow into the foot; **and**
 4. An attestation by a behavioral health provider (i.e., a face-to-face or virtual assessment, with or without psychological questionnaires) reveals no evidence of an inadequately controlled behavioral health condition/issue (e.g., substance use disorder, depression, or psychosis) that would impact perception of pain and/or negatively impact the success of an SCS or contraindicate its placement.
 - B. Permanent implantation of a non-high-frequency dorsal column SCS, when at least a 50% reduction in pain has been demonstrated during a short-term trial of SCS.
- IV. Based upon our criteria and assessment of the peer-reviewed literature, use of a non-high-frequency dorsal column SCS has been medically proven to be effective and, therefore, is considered **medically appropriate** for treatment of patients with chronic, intractable pain secondary to chronic stable angina pectoris or myocardial ischemia, as follows:
 - A. A short-term trial (e.g., greater than 48 hours) of a non-high-frequency dorsal column SCS, when **ALL** the following criteria are met:
 1. Angina pectoris is Canadian Cardiovascular Society (CCS) functional class III or class IV (see Description section below);
 2. An attestation from the patient's treating cardiologist confirms that the individual has coronary artery disease (CAD) **and** is not a suitable candidate for a revascularization procedure;
 3. Optimal medical treatment (OMT) that has failed to adequately improve anginal symptoms, including **all** the following:
 - a. anti-platelet therapy;
 - b. statin and/or other lipid-lowering therapy;
 - c. anti-anginal therapy implemented to pursue a goal heart rate of 60 beats per minute; **and**
 - d. anti-hypertensive therapy as may be indicated to pursue a goal systolic blood pressure (SBP) of less than 140 mmHG and a goal diastolic blood pressure (DBP) of less than 90 mmHG; **and**
 4. An attestation by a behavioral health provider (i.e., a face-to-face or virtual assessment, with or without psychological questionnaires) reveals no evidence of an inadequately controlled behavioral health condition/issue (e.g., substance use disorder, depression, or psychosis) that would impact perception of pain and/or negatively impact the success of an SCS or contraindicate its placement.
 - B. Permanent implantation of a non-high-frequency dorsal column SCS, when there has been a beneficial clinical response during a short-term trial of SCS.
- V. Based upon our criteria and assessment of the peer-reviewed literature, the replacement of an existing high-frequency or non-high-frequency dorsal column SCS and dorsal root ganglion (DRG) stimulator is considered **medically appropriate** when **EITHER** of the following criteria are met:
 - A. The existing stimulator and/or battery/generator is malfunctioning, cannot be repaired, and is no longer under warranty;
 - B. Revision of the electrode percutaneous array(s) or electrode plate/paddle(s) is required.
- VI. Based upon our criteria and assessment of the peer-reviewed literature, a repeat trial of spinal cord or dorsal column stimulator (SCS) following the failure of an initial short-term trial does not improve patient outcomes and, therefore, is considered **not medically necessary** for any indication.
- VII. Based upon our criteria and assessment of the peer-reviewed literature, replacement of a functioning non-high-frequency dorsal column SCS with a high-frequency SCS is considered **not medically necessary**.
- VIII. Based upon our criteria and assessment of the peer-reviewed literature, implantation of a high-frequency or non-high-frequency dorsal column SCS has not been medically proven to be effective and, therefore, is considered investigational for **ANY** other indication, including but not limited to:

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 4 of 16

- A. Post-amputation pain (phantom limb pain);
 - B. Post-herpetic neuralgia;
 - C. Peripheral neuropathy (e.g., chronic intractable pain from diabetic sensory neuropathy);
 - D. Dysesthesias involving the lower extremities secondary to spinal cord injury;
 - E. Abdominal/pelvic visceral pain;
 - F. Chronic cervical or lumbar radiculopathy without prior surgery;
 - G. Chronic cervical, thoracic, or lumbar axial pain without prior spinal surgery;
 - H. Failed cervical and/or thoracic spinal surgery with intractable neuropathic pain in arms(s) or trunk;
 - I. Abdominal pain related to celiac artery compression syndrome;
 - J. Neuropathic pain associated with Multiple Sclerosis.
- IX. Based upon our criteria and assessment of the peer-reviewed literature, implantation of a high-frequency SCS has not been medically proven to be effective and, therefore, is considered **investigational** for **ALL** other indications, including CRPS/RSD.
- X. Based upon our criteria and assessment of the peer-reviewed literature, dorsal root ganglion (DRG) stimulation, including replacement of a dorsal column SCS with a DRG stimulator, has not been medically proven to be effective and, therefore, is considered **investigational** for **ALL** indications, except as noted in Policy Statement V.
- XI. Based upon our criteria and assessment of the peer-reviewed literature, generator modes other than tonic-low and high-frequency (e.g., burst stimulation) has not been medically proven to be effective and, therefore, is considered **investigational**.
- XII. Based upon our criteria and assessment of the peer-reviewed literature, peripheral nerve field stimulation has not been medically proven to be effective and, therefore, is considered **investigational** for treatment of acute or chronic pain conditions, including **ANY** of the following:
- A. FBSS with intractable neuropathic leg pain;
 - B. CRPS/RSD;
 - C. CLI;
 - D. Chronic, stable angina pectoris;
 - E. Post-amputation pain (phantom limb pain);
 - F. Post-herpetic neuralgia;
 - G. Peripheral neuropathy;
 - H. Dysesthesias involving the lower extremities secondary to spinal cord injury.

Refer to Corporate Medical Policy #3.01.02 Psychological Testing

Refer to Corporate Medical Policy #11.01.03 Experimental or Investigational Services

POLICY GUIDELINES

- I. This medical policy does not apply to simple or complex brain, occipital nerve, or peripheral (i.e., cranial nerve, peripheral nerve, autonomic nerve, neuromuscular) neurostimulator pulse generator/transmitter.
- II. A dorsal column stimulator using high-frequency is considered an equally effective alternative to non-high-frequency stimulation only for the treatment of chronic, intractable pain secondary to FBS. A dorsal column SCS capable of using either high-frequency or non-high-frequency stimulation (e.g., dual-mode) is considered an equally effective alternative (when the device uses non-high-frequency stimulation) for the treatment of any of the medically necessary indications/conditions listed above.
- III. The implantation of an SCS is used only as a last resort. Other treatment modalities (pharmacological, surgical, psychological, or physical, if applicable) need to have been tried and failed or have been judged unsuitable or contraindicated. Duration of refractory pain is six (6) months or greater.
- IV. Documentation must reflect an objective measure of a 50% reduction in pain scores with a temporarily implanted electrode, prior to permanent implantation.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 5 of 16

- V. Patients must be carefully screened, evaluated, and diagnosed by a multidisciplinary team, prior to application of these therapies. This evaluation may include a psychological evaluation to exclude any psychiatric or psychosocial history that would negatively influence the outcome of the treatment. *Please to refer to Corporate Medical Policy #3.01.02 Psychological Testing.*

DESCRIPTION

Spinal cord stimulation is a technique for the treatment of chronic pain and involves implantation of electrodes in the epidural space to provide electrical impulses to the spinal cord to inhibit pain transmission to the brain. The procedure initially involves a short-term trial (e. g., greater than 48 hours) of percutaneous temporary spinal cord stimulation to determine whether the spinal cord stimulator device will induce sufficient pain relief to render permanent implantation medically necessary.

Definitions

Complex Regional Pain Syndrome (CRPS), (as defined by the International Association for the Study of Pain [IASP]): a variety of painful conditions following injury which appear regionally having a distal predominance of abnormal findings, exceeding in both magnitude and duration the expected clinical course of the inciting event and often resulting in significant impairment of motor function, and showing variable progression over time. In addition to injury, CRPS can also occur as a result of various medical disorders or illnesses.

Critical Limb Ischemia (CLI): clinical syndrome of ischemic pain at rest and ischemic tissue loss such as non-healing ulcers or gangrene, related to peripheral artery disease (PAD) of the lower limbs. Spinal stimulators may be appropriate for the treatment of intractable rest pain secondary to chronic limb ischemia.

- Ischemic Rest Pain: pain that occurs in the toes or in the area of the metatarsal heads. Occasionally, it occurs in the foot proximal to the metatarsal heads. Elevation of the limb above or at the horizontal position aggravates the pain and pendency, to some degree at least, brings relief. The pain is secondary to severe arterial insufficiency resulting in inadequate perfusion to the distal lower extremity.
- Refer to Rutherford classification table below.

Dorsal Root Ganglion (DRG) Stimulation: an emerging method of treatment for neuropathic pain. With DRG stimulation, leads are placed percutaneously into the epidural space under fluoroscopic guidance directly over the targeted dorsal root ganglion within the lumbar or sacral region of the spine. The procedure initially involves a short-term trial (i.e., greater than 48 hours) using an external pulse generator; upon success of the initial a permanent pulse generator may then be implanted.

- At this time, the evidence in the peer-reviewed scientific literature is insufficient to support long-term safety and efficacy. The use of this technology for treatment of pain conditions remains under investigation.

Failed Back Surgery Syndrome (FBSS): lumbar spinal pain of unknown origin despite surgical intervention or appearing after surgical intervention for spinal pain originally in the same spinal region. Procedures/surgery that do not encroach into the spinal canal (e.g., interspinous/interlaminar/facet distraction, kyphoplasty/vertebroplasty surgery, etc.) are not considered surgical interventions associated with FBSS.

High-Frequency Spinal Cord Stimulation (HF-SCS), (also referred to as kilohertz frequency spinal cord stimulation or HF10): a type of spinal cord stimulation (SCS) providing a higher frequency than traditional spinal cord stimulator systems. The HF10 SCS uses low-amplitude, high-frequency, and short-duration pulses. HF10 SCS does not generate paresthesia and operates at a frequency of 10,000 Hz to provide pain relief in comparison to traditional spinal cord stimulation systems, which operate at a frequency in the range of 40-60 Hz and do generate paresthesia. As an alternative to traditional dorsal spinal column stimulation, HF10 SCS is proven safe and effective for treatment of chronic, intractable low back and leg pain in patients with failed back surgery syndrome (FBSS).

Peripheral Nerve Field Stimulation: a technology that involves placement of electrodes subcutaneously within an area of maximal pain, with the objective of stimulating a region of affected nerves to reduce pain. Depending on the targeted nerve, leads may be placed percutaneously just under the skin or via an open approach for larger deeper peripheral nerves.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 6 of 16

- The use of this technology (used alone or in combination with spinal cord stimulation) for treatment of pain conditions is under investigation.

Peripheral Nerve Stimulation: involves implantation of electrodes near or on a peripheral nerve to reduce pain.

- The use of this technology (used alone or in combination with spinal cord stimulation) for treatment of pain conditions is under investigation.

Spinal Cord Stimulation (SCS), (also known as dorsal column stimulation or neuromodulation): a reversible therapy applied for neuropathic pain with techniques that include multi-output implanted pulse generator and a choice of electrodes, some of which can be placed percutaneously. The technical goal of this therapy is to achieve stimulation of paresthesia of the dorsal horn of the spinal cord at a subjectively comfortable level, overlapping an individual's topography of pain.

Rutherford Classification (Rutherford et al., 1997), refer to the policy statement for treatment of patients with chronic, intractable pain secondary to chronic critical limb ischemia (CLI):

Grade	Category	Clinical Description	Objective Criteria
0	0	Asymptomatic- no hemodynamically significant occlusive disease	Normal treadmill or reactive hyperemia test
	1	Mild claudication	Completes treadmill exercise; AP after exercise > 50 mmHg, but at least 20 mmHg lower than resting value
I	2	Moderate claudication	Between categories 1 and 3
	3	Severe claudication	Cannot complete standard treadmill exercise and AP after exercise < 50 mmHg
II	4	Ischemic rest pain	Resting AP < 40 mmHg, flat or barely pulsatile ankle or metatarsal PVR; TP < 30 mmHg
III	5	Minor tissue loss non-healing ulcer, focal gangrene with diffuse pedal ischemia	Resting AP < 60 mmHg, ankle or metatarsal PVR flat or barely pulsatile; TP < 40 mmHg
	6	Major tissue loss- extending above TM level, functional foot no longer salvageable	Same as category 5

AP: ankle pressure; PVR: pulse volume recording; TM: trans metatarsal; TP: toe pressure

Canadian Cardiovascular Society (CCS) Functional Classifications, refer to the policy statement for treatment of patients with chronic, intractable pain secondary to chronic stable angina pectoris or myocardial ischemia.

Grade	Clinical Description
I.	Ordinary physical activity does not cause angina, such as walking and climbing stairs. Angina occurs with strenuous, rapid, or prolonged exertion at work or recreation.
II.	Slight limitation of ordinary activity. Walking or climbing stairs rapidly, walking uphill, walking or stair-climbing after meals, in cold, in wind, or under emotional stress, or only during the few hours after awakening. Walking more than two blocks on the level and climbing more than one flight of ordinary stairs at a normal pace and in normal conditions.
III.	Marked limitation of ordinary physical activity. Walking one to two blocks on the level and climbing one flight in normal conditions and at a normal pace.
IV.	Inability to carry on any physical activity without discomfort—anginal syndrome may be present at rest.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 7 of 16

RATIONALE

North American Spine Society's (NASS) coverage recommendations outline contradictions to for SCS, including repeating an SCS trial in the same region with the same or similar device for a previously failed trial existing and for untreated drug addiction or poorly controlled psychiatric/psychological disorders and pregnancy. Demand-type cardiac pacemakers are considered a relative contraindication, as it is necessary to interrogate the pacemaker device during the trial to ensure that there is no interference between the SCS pulse generator and the pacemaker.

Traditional Stimulation

Totally implantable dorsal column SCS systems are regulated by the U.S. Food and Drug Administration (FDA) as class III pre-market-approval (PMA) devices. Several devices have received FDA PMA approval. Examples of these devices include, but are not limited to, the Precision Spinal Cord Stimulator System and the Genesis IPG System. Systems with external transmitters are regulated by the FDA as Class II Section 510(k) devices. The FDA granted Section 510(k) approval for Advanced Neuromodulation Systems to market its Renew SCS, to Medtronic to market its Spinal Cord and Peripheral Nerve Stimulation Systems, and to Micronet Medical, Inc. to market its Axxess Spinal Cord Stimulation Lead. St. Jude Medical has also received FDA approval for its Protege MRI spinal cord stimulation system.

There is sufficient evidence in the peer-reviewed literature to permit conclusions that the technology provides significant and sustained relief of pain with minimal side effects in appropriately selected patients with chronic, nonmalignant pain. Studies investigating the effectiveness of spinal cord stimulation as a treatment for patients with chronic back/extremity pain report successful management of pain, a substantial decrease in narcotic use, and an improvement in the quality of life. Studies support the use of spinal cord stimulation for patients with CRPS in the upper extremities through outcomes that demonstrate reduction in pain intensity and increased quality of life (e.g., Harke et al., 2005; Kemler et al., 2006; Kumar et al., 2011; Geurts et al., 2013).

One essential step toward the effective use of SCS devices in potential patients is a trial of the system through percutaneous lead placement. This trial will determine the effectiveness in relieving pain (greater than 50% pain relief) and improving the quality of life in patients with refractory neuropathic pain.

Literature exists to support the value of a presurgical psychological evaluation, to identify factors that may adversely impact functional outcomes after spinal cord stimulation (Doleys, 2006; Heckler et al., 2007; Celestin et al., 2009; NASS, 2017).

There is evidence to favor SCS over standard conservative treatment to improve limb salvage and clinical situations in patients with inoperable CLI (Ubbink et al., 2013 and Conte et al., 2019).

Studies found that SCS improved both the quality of life and cardiac parameters of patients with refractory angina pectoris (Pan et al., 2017).

SCS has also been investigated as a treatment for pain associated with cervical trauma or disc herniation, however further research is needed on the use of SCS treat patients with cervical trauma/disc herniation presenting with arm pain, neck pain, and/or cervicogenic headache.

In 2023, authors of a large (n=7560) real-world, propensity-matched, comparative effectiveness research study reported findings and conclusion that permanent SCS placement was not associated with a meaningful reduction in use of pharmacologic (including opioids) or nonpharmacologic interventions used for chronic pain at 2 years (Dhruva et al., 2023). Although the authors noted six limitations of their study (e.g., observational study design, inability to distinguish the benefit from SCS but required mediations/procedures for other areas of pain, functional measures such as quality of life were not assessed), the authors concluded the results suggest a lack of clinical benefit for most patients and possible harm to some, and suggest that there may be opportunities to redeploy the high—and increasing—use and spending associated with SCS toward more evidence-based interventions for chronic pain relief.

High-Frequency Stimulation

As an alternative to traditional dorsal spinal column stimulation, HF 10 spinal cord stimulation is proven safe and effective for the treatment of chronic, intractable low-back and leg pain in patients with FBSS.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 8 of 16

Nevro (Menlo Park, Calif) gained FDA approval in May 2015 for its Senza SCS system, intended for chronic pain treatment. The device administers the company's HF10 therapy in the trunk and/or limbs, which treats unilateral or bilateral pain related to FBSS, intractable low-back pain, and leg pain. The therapy is the only SCS therapy that is FDA-indicated to alleviate pain without paresthesia (a constant tingling sensation associated with traditional spinal cord stimulation techniques).

In July 2021, the FDA expanded the PMA indications for Nevro's Senza SCS System when programmed to a frequency of 10k Hz to aid in the management of chronic intractable pain of the lower limb(s) associated with diabetic neuropathy. A six-month RCT (Peterson et al., 2021) of 216 patients with painful diabetic neuropathy demonstrated significant improvement in mean VAS score, neurologic examination, and health-related quality of life scores in the SCS group compared to conventional medical management alone. Longer-term studies are needed to confirm durability of effect.

Burst Stimulation

In October 2016, the FDA approved BurstDR stimulation (St. Jude Medical), a clinician programmer application that provides intermittent "burst" stimulation for patients rather than at a constant ("tonic") rate. Burst stimulation is proposed to relieve pain with fewer paresthesia. The burst stimulation device works in conjunction with standard SCS devices. In February 2023, the FDA expanded the Indication for Use for Abbott's Prodigy, Proclaim, and Proclaim XR SCS Systems to include treatment of diabetic peripheral neuropathy of the lower extremities through a series of consistent stimulation pulses, called the tonic stimulation mode.

The SUNBURST (Success Using Neuromodulation with BURST) trial (Deer et al., 2018) was designed to assess the effects of Burst stimulation from St Jude Medical and enrolled 100 patients from 20 centers across the United States randomized to either receive tonic stimulation prior to Burst stimulation, or to receive Burst stimulation prior to tonic stimulation. Forty-five patients were randomized to spinal cord stimulation then burst, and the remaining 55 were randomized to burst then spinal cord stimulation. At the end of the second crossover period, patients were allowed to choose the stimulation mode they preferred and were followed for one year. The study met its primary endpoint of non-inferiority and achieved statistical significance for its pre-specified secondary endpoint of superiority demonstrating patients receiving St. Jude Medical's Burst stimulation achieved superior pain relief and greater treatment success when compared to patients receiving traditional SCS. The estimated difference in the overall visual analog scale score between burst and spinal cord stimulation was -5.1 mm (95% upper CI, -1.14 mm), demonstrating noninferiority ($p < 0.001$) and superiority ($p < 0.017$). The proportion of patients with a decrease in visual analog scale score of 30% or more was 60% (60/100) during burst stimulation and 51% (51/100) during spinal cord stimulation. The proportion of patients whose global impression was minimally improved, moderately improved, or very much improved was approximately 74% in both groups. The authors reported that the programming parameters were not standardized at the beginning of the study but a more standardized approach with lower amplitudes was implemented as the trial was ongoing. Trial limitations included the crossover design, which limits comparison of pain over longer periods of time, lack of blinding, and variable burst programming parameters.

Dorsal Root Ganglion Stimulation

DRG stimulation is an emerging method of treatment for neuropathic pain. With DRG stimulation, leads are placed percutaneously into the epidural space, under fluoroscopic guidance, directly over the targeted dorsal root ganglion within the lumbar or sacral region of the spine. Similar to spinal cord stimulation, a short-term trial (i.e., greater than 48 hours) is recommended, using an external pulse generator; upon success of the trial, a permanent pulse generator may then be implanted. At this time, the evidence in the peer-reviewed scientific literature is insufficient to support long-term safety and efficacy. The use of this technology for treatment of pain conditions remains under investigation.

Peripheral Nerve Field Stimulation (PNFS)

Peripheral nerve field stimulation (also known as peripheral subcutaneous field stimulation) is a form of neuromodulation intended to treat chronic neuropathic pain by placing leads subcutaneously within the area of maximal pain. This technique is different from peripheral nerve stimulation (PNS) whereby specific, visible and identifiable peripheral nerves are target. PNFS is being investigated for low back pain, neck and shoulder pain, inguinal and pelvic pain, thoracic pain, abdominal pain, fibromyalgia, and postherpetic neuralgia. The evidence is insufficient to determine that the technology

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 9 of 16

results in an improvement in the net health outcome (Eldabe et al., 2019; Mironer et al., 2011; McRoberts et al., 2013; Kloimstein et al., 2014; Verrills et al., 2014 and 2011).

In 2013, without more recent update, NICE issued guidance on peripheral subcutaneous field stimulation for chronic low back pain, which stated “Current evidence on the efficacy of peripheral nerve-field stimulation for chronic low back pain is limited in both quantity and quality, and duration of follow-up is limited. Evidence on safety is also limited and there is a risk of complications from any implanted device.”

Closed-Loop Spinal Cord Stimulation

A novel spinal cord stimulation system, the Evoke Spinal Cord Stimulation (SCS) System, provides the first in vivo, real-time, continuous objective measure of spinal cord activation in response to therapy via recorded evoked compound action potentials (ECAPs) in patients during daily use. The Evoke SCS System is an implanted, rechargeable spinal cord stimulation system intended to treat long-term (chronic) pain in the trunk or limbs that are difficult to manage (intractable) and received Federal and Drug Administration (FDA) approval on February 28, 2022. The system is designed to operate in either of two modes: an evoked compound action potential (ECAP) controlled closed-loop stimulation mode or an open-loop (fixed output) stimulation mode. The open-loop stimulation mode is equivalent to that of traditional SCS, and the closed-loop purportedly can provide real-time measurement and automatic adjustment of the strength of the stimulation based on the reading, recording, and response to the ECAP.

Mekhail et al. (2020 and 2022) designed a study to examine pain relief and the extent of spinal cord activation with evoked compound action potentials (ECAPs)-controlled closed-loop versus fixed-output, open-loop spinal cord stimulation for the treatment of chronic back and leg pain. This study is the first to record in-vivo human spinal cord electrophysiology in both stimulation modes and reported that more closed-loop group patients as responders ($\geq 50\%$ reduction) in overall pain 53 of 67 [79.1%] versus 36 of 67 [53.7%] in the open-loop group.

Brooker et al. (2021) reported research findings from the Avalon study, which was also designed to investigate the use of the first closed-loop SCS system in patients with chronic pain. This is a prospective, multicenter, single-arm study where 50 patients were enrolled and followed at one, three, six, 12, 15, 18, 21, and 24 months post permanent implantation of the Evoke SCS System. Although the reported 24-month results support the 12-month results of both this Avalon study and the Evoke study, the study has limitations, and the technology remains under investigation.

CODES

- Eligibility for reimbursement is based upon the benefits set forth in the member’s subscriber contract.
- **CODES MAY NOT BE COVERED UNDER ALL CIRCUMSTANCES. PLEASE READ THE POLICY AND GUIDELINES STATEMENTS CAREFULLY.**
- Codes may not be all inclusive as the AMA and CMS code updates may occur more frequently than policy updates.
- Code Key: Experimental/Investigational = (E/I), Not medically necessary/ appropriate = (NMN).

CPT Codes

Code	Description
0784T	Insertion or replacement of percutaneous electrode array, spinal, with integrated neurostimulator, including imaging guidance, when performed
0785T	Revision or removal of neurostimulator electrode array, spinal, with integrated neurostimulator
63650	Percutaneous implantation of neurostimulator electrode array; epidural
63655	Laminectomy for implantation neurostimulator electrode, plate/paddle; epidural

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION**Policy Number: 7.01.51****Page: 10 of 16**

Code	Description
63661	Removal of spinal neurostimulator electrode percutaneous array(s), including fluoroscopy, when performed
63662	Removal of spinal neurostimulator electrode plate/paddle(s) placed via laminotomy or laminectomy, including fluoroscopy when performed
63663	Revision including replacement, when performed, of spinal neurostimulator electrode percutaneous array(s) including fluoroscopy, when performed
63664	Revision including replacement, when performed, of spinal neurostimulator electrode plate/paddle(s) placed via laminotomy or laminectomy, including fluoroscopy, when performed
63685	Insertion or replacement of spinal neurostimulator pulse generator or receiver requiring pocket creation and connection between electrode array and pulse generator or receiver
63688	Revision or removal of implanted spinal neurostimulator pulse generator or receiver

*Copyright © 2024 American Medical Association, Chicago, IL***HCPCS Codes**

Code	Description
C1820	Generator, neurostimulator (implantable), with rechargeable battery and charging system
C1822	Generator, neurostimulator (implantable), high frequency, with rechargeable battery and charging system
C1827	Generator, neurostimulator (implantable), nonrechargeable, with implantable stimulation lead and external paired stimulation controller
L8679	Implantable neurostimulator pulse generator, any type
L8680	Implantable neurostimulator electrode, each
L8681	Patient programmer (external) for use with implantable programmable neurostimulator pulse generator, replacement only
L8682	Implantable neurostimulator radiofrequency receiver
L8683	Radiofrequency transmitter (external) for use with implantable neurostimulator radiofrequency receiver
L8685	Implantable neurostimulator pulse generator, single array, rechargeable, includes extension
L8686	Implantable neurostimulator pulse generator, single array, non-rechargeable, includes extension
L8687	Implantable neurostimulator pulse generator, dual array, rechargeable, includes extension
L8688	Implantable neurostimulator pulse generator, dual array, non-rechargeable, includes extension
L8689	External recharging system for battery (internal) for use with implantable neurostimulator, replacement only
L8695	External recharging system for battery (external) for use with implantable neurostimulator, replacement only

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 11 of 16

ICD10 Codes

Code	Description
Multiple diagnosis codes	

REFERENCES

- *Al-Kaisy A, et al. Sustained effectiveness of 10kHz high-frequency spinal cord stimulation for patients with chronic, low back pain: 24-month results of a prospective multicenter study. Pain Med 2014 Mar;15(3):347-54.
- *Al-Kaisy A, et al. 10 kHz high-frequency spinal cord stimulation for chronic axial low back pain in patients with no history of spinal surgery: a preliminary, prospective, open-label and proof-of-concept study. Neuromodulation 2017 Jan;20(1):63-70.
- *Amann W, et al. Spinal cord stimulation in the treatment of nonreconstructable stable critical leg ischemia: results of the European Peripheral Vascular Disease Outcome Study (SCS-EPOS). Eur J Vasc Endovasc Surg 2003 Sep;26(3):280-6.
- *American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. Practice guidelines for chronic pain management. Anesthesiol 2010 Apr;112(4):1-24.
- *Andrell P, et al. Long-term effects of spinal cord stimulation on angina symptoms and quality of life in patients with refractory angina pectoris- results from the European Angina Registry Link Study (EARL). Heart 2010 Jul;96(14):1132-6.
- *Blonde L, et al. American Association of Clinical Endocrinology clinical practice guideline: developing a diabetes mellitus comprehensive care plan-2022 Update. Endocr Pract 2022 Oct;28(10):923-1049.
- *Bolash R, et al. Wireless high-frequency spinal cord stimulation (10kHz) compared with multiwaveform low-frequency spinal cord stimulation in the management of chronic pain in failed back surgery syndrome subjects: preliminary results of a multicenter, prospective randomized controlled study. Pain Med 2019 Oct 1;20(10):1971-1979.
- *Bondesson S, et al. Comparison of patients undergoing enhanced external counterpulsation and spinal cord stimulation for refractory angina pectoris. Coron Artery Dis 2008 Dec;19(8):627-34.
- *Borjesson M, et al. Spinal cord stimulation in severe angina pectoris- a systematic review based on the Swedish Council on technology assessment in health care report on long-standing pain. Pain 2008 Dec;140(3):501-8.
- *Brooker C, et al. ECAP-Controlled closed-loop spinal cord stimulation efficacy and opioid reduction over 24-months: final results of the prospective, multicenter, open-label Avalon study. Pain Practice 2021;21(6):680-691.
- *Buchser E, et al. Spinal cord stimulation for the management of refractory angina pectoris. J Pain Symptom Manag 2006 Apr;31(4 Suppl):S36-42.
- *Canadian Cardiovascular Society. Guidelines and clinical practice update library. Angina pectoris, a CCS Grading Scale. 1976. [<https://ccs.ca/guidelines-and-clinical-practice-update-library/>] accessed 05/15/24.
- Conger A, et al. The effectiveness of spinal cord stimulation for the treatment of axial low back pain: a systematic review with narrative synthesis. Pain Med 2020 Nov 1;21(11):2699-2712.
- *Conte MS, et al. Global vascular guidelines on the management of chronic limb-threatening ischemia. J Vasc Surg 2019 Jun;69(6S):3S-125S.e40.
- *Celestin J, et al. Pretreatment psychosocial variables as predictors of outcomes following lumbar surgery and spinal cord stimulation: a systematic review and literature synthesis. Pain Med 2009 May-Jun;10(4):639-53.
- Deer T, et al. Safety analysis of dorsal root ganglion stimulation in the treatment of chronic pain. Neuromodulation 2020 Feb;23(2):239-244.
- *Deer TR, et al. Spinal cord stimulation for refractory angina pectoris and peripheral vascular disease. Pain Physician 2006 Oct;9(4):347-52.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 12 of 16

*Deer TR, et al. A prospective study of dorsal root ganglion stimulation for relief of chronic pain. Neuromodulation 2013 Jan-Feb;16(1):67-71.

Deer TR, et al. The Neuromodulation Appropriateness Consensus Committee on best practices for dorsal root ganglion stimulation. Neuromodulation 2019 Jan;22(1):1-35.

*deVos CC, et al. Burst spinal cord stimulation evaluated in patients with failed back pain surgery syndrome and painful diabetic neuropathy. Neuromodulation 2014 Feb;17(2):152-9.

*Dhruva SS, Murillo J, Ameli O, et al. Long-term outcomes in use of opioids, nonpharmacologic pain interventions, and total costs of spinal cord stimulators compared with conventional medical therapy for chronic pain. JAMA Neurol 2023 Jan;80(1):18-29. PMID: 36441532

*Di Pede F, et al. Immediate and long-term clinical outcome after spinal cord stimulation for refractory stable angina pectoris. Am J Cardiol 2003 Apr 15;91(8):951-5.

*Doleys DM. Psychological factors in spinal cord stimulation therapy: brief review and discussion. Neurosurg Focus 2006 Dec 15;21(6):E1.

*Domingues L, et al. Is a combined programme of manual therapy and exercise more effective than usual care in patients with non-specific chronic neck pain? A randomized controlled trial. Clin Rehabil 2019 Dec;33(12):1908-1918.

Duarte RV, et al. Spinal cord stimulation for the management of painful diabetic neuropathy: a systematic review and meta-analysis of individual patient and aggregate data. Pain 2021 Mar 9. [Online ahead of print].

*Eddicks S, et al. Thoracic spinal cord stimulation improves functional status and relieves symptoms in patients with refractory angina pectoris: the first placebo-controlled study. Heart 2007 May;93(5):585-90.

Eldabe S, et al. Does a screening trial for spinal cord stimulation in patients with chronic pain of neuropathic origin have clinical utility and cost-effectiveness (TRIAL-STIM)? A randomized controlled trial. Pain 2020 Dec;161(12):2820-2829.

*Eldabe S et al. A randomized controlled trial of subcutaneous nerve stimulation for back pain due to failed back surgery syndrome: the SubQStim study. Neuromodulation 2019 Jul;22(5):519-528. PMID 29704437

Food and Drug Administration. Stimwave Technologies Freedom Spinal Cord Stimulator (SCS System). [http://www.accessdata.fda.gov/cdrh_docs/pdf14/K141399.pdf] accessed 05/15/24.

*Frey ME, et al. Spinal cord stimulation for patients with failed back syndrome: A systematic review. Pain Physician 2009 Mar-Apr;12(2):379-97.

*Geurts JW, et al. Spinal cord stimulation for complex regional pain syndrome type I: a prospective cohort study with long-term follow-up. Neuromodulation 2013 Nov-Dec;16(6):523-9.

*Gibbons RJ, et al. ACC/AHA 2002 Guideline update for the management of patients with chronic stable angina-summary article. Circ 2003 Jan;107:149-58.

*Grider JS, et al. Effectiveness of spinal cord stimulation in chronic spinal pain: a systematic review. Pain Physician 2016 Jan;19(1):E33-E54.

*Guzmán Pavón MJ, et al. Comparative effectiveness of manual therapy interventions on pain and pressure pain threshold in patients with myofascial trigger points. Clin J Pain 2022 Dec 1;38(12):749-760.

*Harden RN, et al. Complex regional pain syndrome: practical diagnostic and treatment guidelines, 5th ed. Pain Med 2022 Jun 10;23(Suppl 1):S1-S53.

*Harke H, et al. Spinal cord stimulation in sympathetically maintained complex regional pain syndrome type I with severe disability. A prospective clinical study. Eur J Pain 2005 Aug;9(4):363-73.

*Harney D, et al. Complex regional pain syndrome: the case for spinal cord stimulation (a brief review). Injury 2005 Mar;36(3):357-62.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 13 of 16

Head J, et al. Waves of pain relief: a systematic review of clinical trials in spinal cord stimulation waveforms for the treatment of chronic neuropathic low back and leg pain. World Neurosurg 2019 Nov;131:264-274.e3.

*Heckler DR, et al. Presurgical Behavioral Medicine Evaluation (PBME) for implantable devices for pain management: a 1-year prospective study. Pain Pract 2007 Jun;7(2):110-22.

*Henning H, et al. Spinal cord stimulation in postherpetic neuralgia and in acute herpes zoster pain. Anes Analges 2002 Mar;94(3):694-700.

*Hidalgo B, et al. The efficacy of manual therapy and exercise for treating non-specific neck pain: A systematic review. J Back Musculoskelet Rehabil 2017 Nov 6;30(6):1149 - 1169.

Hoelzer BC, et al. Indirect comparison of 10 kHz spinal cord stimulation (SCS) versus traditional low-frequency SCS for the treatment of painful diabetic neuropathy: a systematic review of randomized controlled trials. Biomedicines 2022;10(10):2630.

Hunter CW, et al. BURST(able): A retrospective, multicenter study examining the impact of spinal cord stimulation with burst on pain and opioid consumption in the setting of salvage treatment and "upgrade". Pain Physician 2020 Nov;23(6):E643-E658.

Huygen FJPM, et al. Effectiveness and safety of dorsal root ganglion stimulation for the treatment of chronic pain: a pooled analysis. Neuromodulation 2020 Feb;23(2):213-221.

*Kapural L, et al. Novel 10-kHz high frequency therapy (HF10 Therapy) is superior to traditional low-frequency spinal cord stimulation for the treatment of chronic back and leg pain: The SENZA-RCT randomized controlled trial. Anesthesiology 2015;123(4):851-860.

*Kapural L, et al. Comparison of 10-kHz high-frequency and traditional low-frequency spinal cord stimulation for the treatment of chronic back and leg pain: 24-month results from a multicenter, randomized, controlled pivotal trial. Neurosurgery 2016 Nov;79(5):667-677.

*Kapural L, et al. Clinical evidence for spinal cord stimulation for failed back surgery syndrome (FBSS): systematic review. Spine (Phila Pa 1976) 2017 Jul 15;42 Suppl 14:S61-S66.

*Kapural L, et al. Technical Aspects of Spinal Cord Stimulation for Managing Chronic Visceral Abdominal Pain: The Results from the National Survey. Pain Medicine 2010 May;11(5):685-691

*Kemler MA, et al. Spinal cord stimulation in patients with chronic reflex sympathetic dystrophy. NEJM 2000 Aug 31;343(9):618-24.

*Kemler MA, et al. Impact of spinal cord stimulation on sensory characteristics in complex regional pain syndrome type I: A randomized trial. Anesthesiol 2001 Jul;95(1):72-80.

*Kemler MA, et al. Spinal cord stimulation for chronic reflex sympathetic dystrophy- five-year follow-up. N Engl J Med 2006 Jun 1;354(22):2394-6.

* Kloimstein H, et al. Peripheral nerve field stimulation (PNFS) in chronic low back pain: a prospective multicenter study. Neuromodulation 2014 Feb;17(2):180-7. PMID 24320718.

*Klomp HM, et al. Spinal cord stimulation is not cost-effective for non-surgical management of critical limb ischaemia. Eur J Vasc Endovas Surg 2006 May;31(5):500-8.

*Klomp HM, et al. What is the evidence on efficacy of spinal cord stimulation in (subgroups of) patients with critical limb ischemia? Ann Vasc Surg 2009 May-Jun;23(3):355-63.

*Kriek N, et al. Comparison of tonic spinal cord stimulation, high-frequency and burst stimulation in patients with complex regional pain syndrome: a double-blind, randomized placebo-controlled trial. BMC Musculoskelet Disord 2015 Aug 25;16:222.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 14 of 16

- *Kumar K, et al. Spinal cord stimulation versus conventional medical management for neuropathic pain: a multicentre randomised controlled trial in patients with failed back surgery syndrome. Pain 2007 Nov;132(1-2):179-88.
- *Kumar K, et al. The effects of spinal cord stimulation in neuropathic pain are sustained: a 24-month follow-up of the prospective randomized controlled multicenter trial of the effectiveness of spinal cord stimulation. Neurosurg 2008 Oct;63(4):762-70.
- *Kumar K, et al. Spinal cord stimulation is effective in management of complex regional pain syndrome I: fact or fiction. Neurosurgery 2011 Sep;69(3):566-78.
- *Lanza GA, et al. Effect of spinal cord stimulation on spontaneous and stress reduced angina and ischemic-like ST-segment depression in patients with cardiac syndrome X. Eur Heart J 2005 Jan 9.
- *Lanza GA, et al. Spinal cord stimulation for the treatment of refractory angina pectoris: a multicenter randomized single-blind study (the SCS-ITA trial). Pain 2011 Jan;152(1):45-52.
- *Liem L, et al. A multicenter, prospective trial to assess the safety and performance of the spinal modulation dorsal root ganglion neurostimulator system in the treatment of chronic pain. Neuromodulation 2013 Sept-Oct;16(5):471-482.
- *Liem L, et al. One-year outcomes of spinal cord stimulation of the dorsal root ganglion in the treatment of chronic neuropathic pain. Neuromodulation 2015 Jan;18(1):41-48.
- Lin I, et al. What does best practice care for musculoskeletal pain look like? Eleven consistent recommendations from high-quality clinical practice guidelines: systematic review. Br J Sports Med 2020 Jan;54(2):79-86. Epub 2019 Mar 2.
- *Mailis-Gagnon A, et al. Spinal cord stimulation for chronic pain. Cochrane Database Syst Review 2004;(3): CD003783.
- *Manca A, et al. Quality of life, resource consumption and costs of spinal cord stimulation versus conventional medical management in neuropathic pain patients with failed back surgery syndrome (PROCESS trial). Eur J Pain 2008 Nov;12(8):1047-58.
- *McNab D, et al. An open label, single-centre, randomized trial of spinal cord stimulation vs. percutaneous myocardial laser revascularization in patients with refractory angina pectoris: the SPiRiT trial. Eur Heart J 2006 May;27(9):1048-53.
- *McRoberts WP, et al. Peripheral nerve field stimulation for the management of localized chronic intractable back pain: results from a randomized controlled study. Neuromodulation 2013;16(6):565-574. PMID 23577773
- *Mekhail NA, et al. Retrospective review of 707 cases of spinal cord stimulation: indications and complications. Pain Pract 2011 Mar;11(12):148-53.
- *Mekhail NA, et al. Long-term safety, and efficacy of closed-loop spinal cord stimulation to treat chronic back and leg pain (Evoke): a double-blind, randomised, controlled trial. Lancet Neurol 2020(19):123-34.
- *Mekhail NA, et al. Durability of clinical and quality-of-life outcomes of closed-loop spinal cord stimulation for chronic back and leg pain. JAMA Neurol 2022 Mar;79(3):251-260.
- *Mironer YE, et al. Prospective, two-part study of the interaction between spinal cord stimulation and peripheral nerve field stimulation in patients with low back pain: development of a new spinal-peripheral neurostimulation method. Neuromodulation 2011;14(2):151-154. PMID 21992203
- *Morgalla MH, et al. Dorsal root ganglion stimulation (DRGS) for the treatment of chronic neuropathic pain: a single-center study with long-term prospective results in 62 cases. Pain Physician 2018 Jul;21(4):E377-E387.
- *National Institute for Health and Care Excellence (NICE). Peripheral nerve-field stimulation for chronic low back pain [IPG451]. 2013 Mar 27 [<https://www.nice.org.uk/guidance/ipg451>] accessed 05/15/24.
- *National Institute of Clinical Excellence (NICE). Spinal cord stimulation for chronic pain of neuropathic or ischaemic origin. TA159. 2008 Oct; rev 2014 Feb [<https://www.nice.org.uk/guidance/ta159>] accessed 05/15/24.
- *North American Spine Society. Coverage policy recommendations. Spinal cord stimulation. 2017 Oct. [<https://www.spine.org/Coverage>] accessed 08/30/24.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 15 of 16

- *North RB, et al. Spinal cord stimulation for chronic pain of spinal origin: a valuable long-term solution. Spine 2002 Nov 15; 27(22):2584-91.
- *North RB, et al. Spinal cord stimulation for axial low back pain: a prospective, controlled trial comparing dual and single percutaneous electrodes. Spine 2005 Jun 15;30(12):1412-8.
- *North RB, et al. Spinal cord stimulation versus reoperation for failed back surgery syndrome: a cost effectiveness and cost utility analysis based on a randomized, controlled trial. Neurosurgery 2007 Aug;61(2):361-8.
- *Pan X, Bao H, Si Y, et al. Spinal cord stimulation for refractory angina pectoris: a systematic review and meta-analysis. Clin J Pain Jun 2017; 33(6): 543-551.
- *Pedrini L, et al. Spinal cord stimulation for lower limb ischemic pain treatment. Interact Cardiovasc Thorac Surg 2007 Aug;6(4):495-500.
- *Perruchoud C, et al. Analgesic efficacy of high-frequency spinal cord stimulation: a randomized double-blind placebo-controlled study. Neuromodulation 2013 July-Aug;16(4):363-369.
- *Petersen EA, et al. Effect of high-frequency (10-kHz) spinal cord stimulation in patients with painful diabetic neuropathy: a randomized clinical trial. JAMA Neurol 2021 Jun 1;78(6):687-698.
- Petersen EA, et al. Durability of High-Frequency 10-kHz Spinal Cord Stimulation for Patients With Painful Diabetic Neuropathy Refractory to Conventional Treatments: 12-Month Results From a Randomized Controlled Trial. Diabetes Care 2022 Jan;45:e3-e6.
- Pop-Busui R, et al. Diagnosis and treatment of painful diabetic peripheral neuropathy. ADA Clinical Compendia 2022 Feb;2022(1):1-32.
- *Riberio DC, et al. Mediators of the effects of exercise and manual therapy for people with knee and hip osteoarthritis: A secondary, exploratory analysis of the MOA trial. Osteoarthr Cartil Open 2024 Jan 9;6(1):1-6. eCollection 2024 Mar.
- *Russo M and Van Buyten JP. 10-kHz high frequency SCS therapy: a clinical summary. Pain Med 2015 May;16(5):934-942.
- *Rutherford RB, et al. Recommended standards for reports dealing with lower extremity ischemia: revised version. J Vasc Surg 1997 Sep;26(3):517-38.
- *Schu S, et al. A prospective, randomized, double-blind, placebo-controlled study to examine the effectiveness of burst spinal cord stimulation patterns for the treatment of failed back surgery syndrome. Neuromodulation 2014 Jul;17(5):443-50.
- *Sears NC, et al. Long-term outcomes of spinal cord stimulation with paddle leads in the treatment of complex regional pain syndrome and failed back surgery syndrome. Neuromodulation 2011 Jul-Aug;14(4):312-8.
- *Sgueglia GA, et al. Long-term follow-up of patients with cardiac syndrome X treated by spinal cord stimulation. Heart 2007 May;93(5):591-7.
- *Simpson EL, et al. Spinal cord stimulation for chronic pain of neuropathic or ischaemic origin: systematic review and economic evaluation. Health Tech Assess 2009 Mar;13(17):iii, ix-x,1-154.
- *Sitzman BT and Provenzano DA, et al. Best practices in spinal cord stimulation. Spine (Phila Pa 1976) 2017 Jul 15;42 Suppl 14:S67-S71.
- *Strand N, et al. Evidence-based clinical guidelines from the American Society of Pain and Neuroscience for the use of implantable peripheral nerve stimulation in the treatment of chronic pain. J Pain Res 2022; 15: 2483-2504. PMID 36039168
- *Taylor RS, et al. Spinal cord stimulation for chronic back pain and leg pain and failed back surgery syndrome: a systemic review and analysis of prognostic factors. Spine 2005 Jan 1;30(1):152-60.

Medical Policy: SPINAL CORD STIMULATION/DORSAL COLUMN STIMULATION

Policy Number: 7.01.51

Page: 16 of 16

*Taylor RS. Spinal cord stimulation in complex regional pain syndrome and refractory neuropathic back and leg pain/failed back surgery syndrome: results of systematic review and meta-analysis. J Pain Symptom Manag 2006 Apr;3(4 Suppl):S13-9.

*Taylor RS, et al. Spinal cord stimulation in the treatment of refractory angina: systematic review and meta-analysis of randomized controlled trials. BMC Cardiovasc Disord 2009 Mar 25;9:13.

*Taylor RS, et al. Predictors of pain relief following spinal cord stimulation in chronic back and leg pain and failed back surgery syndrome: A systematic review and meta-regression analysis. Pain Pract 2014 Jul;14(6):489-505.

Traeger AC, et al. Spinal cord stimulation for low back pain. Cochrane Database Syst Rev. 2023 May 07;3(3):CD014789.

*Tsertsvadze A, et al. Cost-effectiveness of manual therapy for the management of musculoskeletal conditions: a systematic review and narrative synthesis of evidence from randomized controlled trials. J Manipulative Physiol Ther 2014; 37(6): 343-62.

*Ubbink DT, et al. Spinal cord stimulation for non-reconstructable chronic critical leg ischaemia. Cochrane Database Syst Rev 2013 Feb 28;2:CD004001.

*Ubbink DT, et al. Spinal cord stimulation for critical leg ischemia: a review of effectiveness and optimal patient selection. J Pain Symptom Manag 2006 Apr;31(4 Suppl):S30-5.

*Urits I, et al. A comprehensive review of alternative therapies for the management of chronic pain patients: acupuncture, tai chi, osteopathic manipulative medicine, and chiropractic care. Adv Ther 2020;38(1):76 - 89. Epub 2020 Nov 12.

*van Beek M, et al. Sustained treatment effect of spinal cord stimulation in painful diabetic peripheral neuropathy: 24-month follow-up of a prospective two-center randomized controlled trial. Diabetes Care 2015 Sept;38(9):e132-134.

*van Buyten JP, et al. High-frequency spinal cord stimulation for the treatment of chronic back pain patients: results of a prospective multicenter European clinical study. Neuromodulation 2013 Jan-Feb;16(1):59-65.

*Verrills P, et al. Peripheral nerve field stimulation for chronic pain: 100 cases and review of the literature. Pain Med 2011 Sep;12(9):1395-405. PMID 21812906

*Verrills P, et al. Peripheral nerve field stimulation for chronic headache: 60 cases and long-term follow-up. Neuromodulation 2014 Jan;17(1):54-59. PMID 24165152

Yeung AM, et al. Spinal cord stimulation for painful diabetic neuropathy. J Diabetes Sci Technol 2024 Jan;18(1):168 192. Epub 2022 Nov 17.

*Zipes DP, et al. Determining the feasibility of spinal cord neuromodulation for the treatment of chronic systolic heart failure: the DEFEAT-HF study. JACC Heart Fail 2016 Feb;4(2):129-136.

Zuidema X, et al. Long-term evaluation of spinal cord stimulation in patients with painful diabetic polyneuropathy: An eight-to-ten-year prospective cohort study. Neuromodulation 2022 Dec 30:S1094-7159(22)01403-9.

*Key Article

KEY WORDS

Burst stimulation, Dorsal column, Dorsal root ganglion, High-frequency neurostimulation, Neuromodulation, Neurostimulation, Wireless neurostimulation, Closed-loop SCS.

CMS COVERAGE FOR MEDICARE PRODUCT MEMBERS

There is currently a National Coverage Determination (NCD) (160.7), Electrical Nerve Stimulators, for dorsal column stimulators. Please refer to the following NCD website for Medicare Members: [<https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?NCID=240>] accessed 08/30/24.

Based on our review, peripheral nerve field stimulation (PNFS) is not addressed in National or Regional Medicare coverage determinations or policies.